


Phenyl addition made a poison useful for a chemical reaction in catalysis

November 16 2018



Take for example a poisoned apple, it can emit an intoxicating mellow aroma just by sticking a pen and change to be delicious. Credit: Figure adapted with permission from *Organometallics*, 2018, 10.1021/acs.organomet.8b00645. © 2018 American Chemical Society

Scientists from Tokyo University of Agriculture and Technology (TUAT), Japan, have discovered that a catalyst poison, which deactivates homogeneous catalysts, can be "reborn" as an efficient ligand by introduction of a substituent, in chemical reactions. This finding is of service to expand the ligand design in homogeneous catalysts. This

research was published online in the journal *Organometallics*.

A group of substances that completely or partly shut down a catalysis by contamination is widely known as [catalyst](#) poison. Typical examples of catalyst poison are mercury and sulfured compounds for [heterogeneous catalysis](#) and a [chemical substance](#) called dibenzocyclooctatetraene for homogeneous catalysis. These substances deactivate the catalyst by binding to the active site very tightly. The homogenous catalyst poison, however, can be changed as an efficient [ligand](#) by introduction of a substituent.

A research team at TUAT was looking for cyclic diene ligands for a ruthenium (Ru) catalyst to synthesize skipped dienes, which are known to be substructures of many natural products and pharmaceutical molecules. By accident, they came across dibenzocyclooctatetraene, a homogenous catalyst poison, to work as a ligand . After screening, they finally found that phenyl attachment to the catalyst poison was a very efficient ligand. Technically, reaction of b-myrcene (one of the most widespread group of natural products) with methyl acrylate catalyzed by a Ru complex with 1,5-cyclooctadiene (1 mol percent; a typical cyclic diene ligand) produced the linear coupling product only in 23 percent yield at 30 degrees C for 7 hours. However, the 5-phenyldibenzocyclooctatetraene (dibenzocyclooctatetraene was attached to a phenyl group) complex of Ru catalyzed the reaction to give the product in 97 percent yield within 1 hour under the same conditions.

The dibenzocyclooctatetraene tightly binds to the Ru center by its nature. One of the keys for this high catalytic activity probably arises from the steric effect of the phenyl group. In the presence catalysis giving skipped dienes, the formed skipped diene molecules are prone to stay on the catalyst as a stable intermediate but the phenyl group sweeps the product molecule away from the catalyst. As the result, the catalyst can take actions for further catalysis.

"Before a catalytic process, one normally has to remove substances that work as the catalyst poisons not to deactivate the catalyst. They are nuisances in catalyses because they tightly bound to the active site in catalysts. However, the strongly bind nature to the catalyst is an advantage as a ligand in [homogeneous catalysts](#). Our finding let us know a small alteration of a catalyst poison, not only do-not-hampers the catalysis, but brings acceleration of the [catalysis](#) as a ligand of catalyst. Our results are surely service to the further development on the design of tough ligands for catalysts. In other words, a catalyst poison is not born this way, but can be reborn as an efficient ligand," said Masafumi Hirano, a TUAT professor of chemistry and a principal of the study.

More information: Masafumi Hirano et al, Synthesis of and Catalytic Linear Cross-Dimerizations by an Electron-Deficient Cyclic Diene Complex of Ruthenium(0), *Organometallics* (2018). [DOI: 10.1021/acs.organomet.8b00645](#)

Provided by Tokyo University of Agriculture and Technology

Citation: Phenyl addition made a poison useful for a chemical reaction in catalysis (2018, November 16) retrieved 29 June 2024 from <https://phys.org/news/2018-11-phenyl-addition-poison-chemical-reaction.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.