


# Species and environment affect which frogs are infected by parasitic fungus

November 19 2018

---



The parasitic fungus *Batrachochytrium dendrobatidis* causes lethal infections in amphibians. Species and environment affect which frogs are infected by parasitic fungus. This is a green toad. Credit: Uppsala University

An aquatic parasitic fungus causes lethal infections in amphibians and is thought to be one of the reasons for a global decline in toad and frog populations. A new study by researchers from Uppsala University shows a wide variation among different species in the number of infections and

that the surrounding environment has an impact.

The study was performed in Skåne, which is home to twelve [amphibian species](#). The common frog, common toad, moor frog, European fire-bellied toad, European green toad and the natterjack toad were included in the study. The number of infections varied widely among the species. The common toad and common frog had the lowest proportion of infections, while the European green toad, the natterjack toad and the fire-bellied toad had the highest. In addition to the differences in the proportions of infected amphibians across the species, the [infection](#) rate was also affected by the environment in the ponds and surrounding landscape where the frogs and toads live. The proportion of infected individuals increased if the pH value in the [pond](#) was high and decreased if there was forest and many nearby ponds in the surrounding countryside.

Earlier studies have shown that the parasitic fungus (*Batrachochytrium dendrobatidis*) is sensitive to temperature and pH. Because the number of infected individuals was higher in ponds with higher pH values, the results suggest that the fungus is happiest in such environments. Likewise, forests can affect the prevalence of the fungus because it is relatively colder in wooded areas than in more open country. Some [amphibian](#) species avoid colder areas and ponds with specific pH values, which also affects the prevalence.

**More information:** Simon Kärnvemo et al. Effects of host species and environmental factors on the prevalence of *Batrachochytrium dendrobatidis* in northern Europe, *PLOS ONE* (2018). [DOI: 10.1371/journal.pone.0199852](https://doi.org/10.1371/journal.pone.0199852)

Provided by Uppsala University

Citation: Species and environment affect which frogs are infected by parasitic fungus (2018, November 19) retrieved 17 May 2024 from <https://phys.org/news/2018-11-species-environment-affect-frogs-infected.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.