



# Table-top experiment flips current understanding of solutal convection

November 30 2018


Experiments



Direct Numerical Simulations without Dispersion



Direct Numerical Simulations with Dispersion



The convection in the right column is much stronger than that in the left. The experiments (top) show that the convective pattern is coarsened at larger grain size (right), while the classical studies (middle) generally predict the opposite. After considering the effect of mechanical dispersion, the new simulations

results (bottom) match well with the experimental observations. Credit: Liang *et al.*

When Yu "Alex" Liang started graduate school at The University of Texas at Austin, he was tasked with running a straight-forward experiment to collect data on a well-understood phenomenon in fluid mechanics: how density differences influence fluid flow in a porous medium.

It's a scenario that plays out across a wide range of scientific problems. In fact, Liang was planning on applying the experimental data toward a larger project on carbon sequestration. However, the experiment revealed that the physics governing the flow was not as settled as scientists thought.

The findings revealed that the primary driver of this type of fluid flow—called solutal [convection](#)—had been overlooked. What's more, once this driver is accounted for, it completely flips the expected flow outcomes.

"To some degree, it is a game changer," said Liang, who earned his Ph.D. in [petroleum engineering](#) in 2017 and is now working for Hilcorp, a Houston-based energy company. "Our experiments and simulations show that the convective pattern is controlled by a different process than the one previously thought. People will realize that there are much more in-depth theories to be explored on solutal convection in porous media."

The results of Liang's research were published in the journal *Geophysical Researcher Letters* in September. His co-authors include Marc Hesse, an associate professor at the UT Jackson School of Geosciences and the Institute for Computational Engineering and Sciences (ICES), David

DiCarlo, an associate professor at the UT Hildebrand Department of Petroleum and Geosystems Engineering, and Baole Wen, a postdoctoral fellow at the Jackson School and ICES. DiCarlo and Hesse are Liang's Ph.D. advisors.

For decades it has been the scientific consensus that the balance between density-driven currents and diffusion was the primary factor controlling solutal convection in porous media. The thinking went: regions of dense fluid move downward until diffusion eliminates the density difference driving the flow, creating dense fingers that sink toward the bottom. In general, the fingers have to be far enough apart so that diffusion cannot smear them out as the fluid sinks. Therefore, the fingers were generally thought to be further apart in slow flows and closer together in fast flows driven by bigger density differences.

Despite the pattern being well-established in [computer simulations](#), Hesse said that the research team was unable to find experimental results that demonstrate this basic behavior. So they developed a simple table-top rig—a transparent tank filled with glass beads and water—for observing solutal convection in real time. To start the convection process, the water was topped with a layer of methanol and ethylene-glycol—a mixture that is overall less dense than water, but gradually becomes denser and sinks as it undergoes convection with the water at the fluid interface.

The team was expecting the classic narrow finger pattern to emerge in experiments that used larger diameter beads. Instead, the complete opposite pattern emerged. The finger spacing increased with bead size.

"Here's this very basic phenomena, which occurs in all sorts of applications, is a classic example of pattern formation—and you do the experiments and you get literally the opposite of what everybody expects," Hesse said. "This shows that something is totally off in our

basic understanding of this process."

A deeper analysis revealed that the dispersion created by the larger diameter beads had a greater impact on the convective environment than diffusion. While previous theories of solutal convection emphasize diffusion, the spreading of the material in the water in their experiments turned out to be controlled by mechanical dispersion, which leads to the additional mixing of the fluids at the pore scale.

"In other words, what you're flowing through is really important," he said. "Dispersion gets bigger with increasing grain size, and that's why the fingers get wider as you do this with bigger beads."

Based on their observations of the table-top experiment, the researchers were able to replicate their findings with a computational model.

"The key point is that we analyze the effect of dispersion on convection in subsurface based on our [experimental data](#) and then use high-resolution numerical simulations to verify our analysis," said Wen.

Of course, the natural world is much more complex than a tank filled with beads or a simplified model. The researchers said that scientists have many factors to keep in mind when researching complex phenomena that involve solutal convection, such as CO<sub>2</sub> sequestration. But DiCarlo said that these findings show that scientists studying the fundamentals of the process have some new factors to consider.

"The work shows that if one wants to predict how CO<sub>2</sub> dissolves in the subsurface, or similar dissolution process, the study needs to include dispersion in the correct manner," he said. "All previous studies have ignored dispersion."

Hesse added that the experimental results can help add a dose of reality

to computational models that have been getting it wrong for decades.

"If your numerical simulation can't even simulate this simple experiment that I'm doing, what confidence do you have that it will do the right thing in an even more complicated setting," he said.

**More information:** Yu Liang et al, Effect of Dispersion on Solutal Convection in Porous Media, *Geophysical Research Letters* (2018). [DOI: 10.1029/2018GL079849](https://doi.org/10.1029/2018GL079849)

Provided by University of Texas at Austin

Citation: Table-top experiment flips current understanding of solutal convection (2018, November 30) retrieved 9 April 2024 from <https://phys.org/news/2018-11-table-top-flips-current-solutal-convection.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--