

## A more accurate way of resolving spatial patterns in weather could lead to better predictions of climate change

December 5 2018



High-resolution reconstruction of rainfall using the local nonstationary model—rainfall is shown as the annual averaged precipitation rate (mm/h). Credit: KAUST

A divide and conquer approach, combined with some mathematical gymnastics, has helped KAUST researchers Ying Sun and Yuxiao Li to come up with a fast and flexible statistical scheme for improving the accuracy of climate-change modeling.



"Statisticians attempt to avoid unrealistic assumptions in order to estimate environmental processes more accurately," says Li, a Ph.D. student in Sun's research team. "The motivation behind our study was to improve how spatial nonstationarity is accounted for in <u>climate</u> modeling over a large and geographically complex area."

Nonstationarity occurs when the statistics of an observed parameter, such as temperature or rainfall, varies over time or space. In this case, Li and Sun looked at the statistics that describe how measurements at two different locations change spatially. Understanding and accurately characterizing this covariance is critical in predicting weather and in translating or downscaling <u>global climate</u> modeling that is done at very coarse scales to be able to predict local effects at finer scales.

"We can think of spatial covariance as how well observations are correlated with each other in space, or as spatial patterns," explains Li. "For environmental data, the spatial correlation pattern over land is different from that over the ocean. Because the correlation pattern is not exactly the same everywhere, we need to consider how to represent this nonstationarity."

Nonstationarity has been dealt with previously by complex and computationally intensive convolution-based approaches, coupled with approximations based on assumptions of local stationarity at an arbitrary grid size. Sun and Li developed a computationally efficient approach that greatly improves the accuracy of approximation.

"The challenge for nonstationary modeling is the trade-off between accuracy and efficiency," says Li. "We extended the local stationary model, which is efficient but not accurate, to a linearly varying local nonstationary model by dividing up the spatial region and estimating the covariance functions for each subregion. This allows us to describe more complicated nonstationary features, which improves accuracy while still



being fast to compute."

The researchers expect to significantly improve the downscaling of climate modeling through this approach, which avoids unrealistic assumptions and can estimate different types of spatial nonstationarity at a finer spatial scale. They demonstrated the efficiency and improved accuracy of their scheme by simulating precipitation at high-spatial resolution over a large region. The approach also lends itself to the interpolation of environmental processes at unobserved locations.

"Since our method is based on more realistic assumptions, analyses using this <u>approach</u> can help scientists better understand weather and climate," says Li.

**More information:** Yuxiao Li et al. Efficient Estimation of Nonstationary Spatial Covariance Functions with Application to Highresolution Climate Model Emulation, *Statistica Sinica* (2018). DOI: <u>10.5705/ss.202017.0536</u>

## Provided by King Abdullah University of Science and Technology

Citation: A more accurate way of resolving spatial patterns in weather could lead to better predictions of climate change (2018, December 5) retrieved 11 July 2024 from <u>https://phys.org/news/2018-12-accurate-spatial-patterns-weather-climate.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.