


Algorithms to locate centrioles in the cell

December 12 2018



The study of these cylindrical organelles of the cellular cytoskeleton has been carried out with the wings of the fruit fly. Credit: UEx

Investigators from the Neurodegenerative Diseases Research Group at the University of Extremadura are studying signaling mediated by a pathway known as planar cell polarity (PCP), which regulates the coordinated orientation of cells during organogenesis, the process of organ formation in living organisms. This pathway has been highly conserved on the evolutionary scale, and one of its key functions in vertebrates is the regulation of the coordinated positioning of

centrioles/ciliary basal cells inside cells.

This signaling pathway was discovered initially in the fruit fly genus *Drosophila*, although the majority of the pathway components have been retained in humans. It has likewise been observed that certain pathologies such as hydrocephaly, infertility and some kinds of cancers are associated with defective functioning of this signaling.

Under the auspices of the project EPICENTR within the Spanish national research plan, whose objective is to study the planar polarisation of centrioles in epithelial [cells](#), the UEx researchers have now published the first results of their investigation in the journal *Development*. These results are related to the polarised positioning mechanism of centrioles in *Drosophila* and its correlation with actin.

"We have developed a methodology with [new algorithms](#) to analyse the location of the centriole in a model cell. Thanks to this technology, we have been able to discover how the actin cytoskeleton, which is part of the cell scaffolding, is involved in the polarised placement of centrioles in *Drosophila*, just as happens in vertebrates. Nevertheless, the well-known signaling route downstream PCP, which controls the [actin cytoskeleton](#), does not affect the positioning of centrioles in *Drosophila*," explains José María Carvajal, lead researcher on this project. The PCP-dependent factors which affect the location of the centriole in humans and in vertebrates are not found in *Drosophila*. It follows that there are new proteins to be discovered in *Drosophila* which influence the actin and the location of centrioles.

"The advantage is that in this investigation, we provide a technology for image analysis. We have applied new algorithms to analyse where the centrioles are in [epithelial cells](#) using an imaging system, yielding a model that enables us to extract the variety of possible positions for the centriole inside a cell," says Carvajal.

Now, the researchers are studying another cell polarity mechanism in human cells by carrying over the *Drosophila* biological model to vertebrates. They will describe its effects on the respiratory system, in the so-called ciliopathies, illnesses arising out of malfunctioning of the cilia inside the cell.

More information: Sergio Garrido-Jimenez et al, Centriole planar polarity assessment in *Drosophila* wings, *Development* (2018). [DOI: 10.1242/dev.169326](https://doi.org/10.1242/dev.169326)

Provided by University of Extremadura

Citation: Algorithms to locate centrioles in the cell (2018, December 12) retrieved 18 April 2024 from <https://phys.org/news/2018-12-algorithms-centrioles-cell.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.