


Biggest mass extinction caused by global warming leaving ocean animals gasping for breath

December 6 2018



This illustration shows the percentage of marine animals that went extinct at the end of the Permian era by latitude, from the model (black line) and from the fossil record (blue dots). A greater percentage of marine animals survived in the tropics than at the poles. The color of the water shows the temperature change, with red being most severe warming and yellow less warming. At the top is the supercontinent Pangaea, with massive volcanic eruptions emitting carbon dioxide. The images below the line represent some of the 96 percent of marine species that died during the event. Includes fossil drawings by Ernst Haeckel/Wikimedia; Blue crab photo by Wendy Kaveney/Flickr; Atlantic cod

photo by Hans-Petter Fjeld/Wikimedia; Chambered nautilus photo by ©2010 John White/CalPhotos. Credit: Justin Penn and Curtis Deutsch/University of Washington

The largest extinction in Earth's history marked the end of the Permian period, some 252 million years ago. Long before dinosaurs, our planet was populated with plants and animals that were mostly obliterated after a series of massive volcanic eruptions in Siberia.

Fossils in ancient seafloor rocks display a thriving and diverse marine ecosystem, then a swath of corpses. Some 96 percent of marine species were wiped out during the "Great Dying," followed by millions of years when life had to multiply and diversify once more.

What has been debated until now is exactly what made the oceans inhospitable to life—the high acidity of the water, metal and sulfide poisoning, a complete lack of oxygen, or simply higher temperatures.

New research from the University of Washington and Stanford University combines models of ocean conditions and animal metabolism with published lab data and paleoceanographic records to show that the Permian mass extinction in the oceans was caused by [global warming](#) that left animals unable to breathe. As temperatures rose and the metabolism of marine animals sped up, the warmer waters could not hold enough oxygen for them to survive.

The study is published in the Dec. 7 issue of *Science*.

"This is the first time that we have made a mechanistic prediction about what caused the extinction that can be directly tested with the fossil record, which then allows us to make predictions about the causes of

extinction in the future," said first author Justin Penn, a UW doctoral student in oceanography.

Researchers ran a climate model with Earth's configuration during the Permian, when the land masses were combined in the supercontinent of Pangaea. Before ongoing volcanic eruptions in Siberia created a greenhouse-gas planet, oceans had temperatures and [oxygen levels](#) similar to today's. The researchers then raised greenhouse gases in the model to the level required to make tropical ocean temperatures at the surface some 10 degrees Celsius (20 degrees Fahrenheit) higher, matching conditions at that time.

The model reproduces the resulting dramatic changes in the oceans. Oceans lost about 80 percent of their oxygen. About half the oceans' seafloor, mostly at deeper depths, became completely oxygen-free.

To analyze the effects on marine species, the researchers considered the varying oxygen and temperature sensitivities of 61 modern [marine species](#)—including crustaceans, fish, shellfish, corals and sharks—using published lab measurements. The tolerance of modern animals to high temperature and low oxygen is expected to be similar to Permian animals because they had evolved under similar environmental conditions. The researchers then combined the species' traits with the paleoclimate simulations to predict the geography of the extinction.

"Very few [marine organisms](#) stayed in the same habitats they were living in—it was either flee or perish," said second author Curtis Deutsch, a UW associate professor of oceanography.



This roughly 1.5-foot slab of rock from southern China shows the Permian-Triassic boundary. The bottom section is pre-extinction limestone. The upper section is microbial limestone deposited after the extinction. Credit: Jonathan Payne/Stanford University

The model shows the hardest hit were organisms most sensitive to oxygen found far from the tropics. Many species that lived in the tropics also went extinct in the model, but it predicts that high-latitude species, especially those with high oxygen demands, were nearly completely wiped out.

To test this prediction, co-authors Jonathan Payne and Erik Sperling at Stanford analyzed late-Permian fossil distributions from the Paleooceanography Database, a virtual archive of published fossil collections. The fossil record shows where species were before the extinction, and which were wiped out completely or restricted to a fraction of their former habitat.

The [fossil record](#) confirms that species far from the equator suffered most during the event.

"The signature of that kill mechanism, climate warming and oxygen loss, is this geographic pattern that's predicted by the model and then discovered in the fossils," Penn said. "The agreement between the two indicates this mechanism of climate warming and oxygen loss was a primary cause of the extinction."

The study builds on previous work led by Deutsch showing that as oceans warm, marine animals' metabolism speeds up, meaning they require more oxygen, while warmer water holds less. That earlier study shows how warmer oceans push animals away from the tropics.

The new study combines the changing ocean conditions with various animals' metabolic needs at different temperatures. Results show that the most severe effects of oxygen deprivation are for species living near the poles.

"Since tropical organisms' metabolisms were already adapted to fairly warm, lower-oxygen conditions, they could move away from the tropics and find the same conditions somewhere else," Deutsch said. "But if an organism was adapted for a cold, oxygen-rich environment, then those conditions ceased to exist in the shallow oceans."

The so-called "dead zones" that are completely devoid of oxygen were mostly below depths where species were living, and played a smaller role in the survival rates." At the end of the day, it turned out that the size of the dead zones really doesn't seem to be the key thing for the extinction," Deutsch said. "We often think about anoxia, the complete lack of oxygen, as the condition you need to get widespread uninhabitability. But when you look at the tolerance for low oxygen, most organisms can be excluded from seawater at oxygen levels that aren't anywhere close to

anoxic."

Warming leading to insufficient [oxygen](#) explains more than half of the marine diversity losses. The authors say that other changes, such as acidification or shifts in the productivity of photosynthetic organisms, likely acted as additional causes.

The situation in the late Permian—increasing greenhouse gases in the atmosphere that create warmer temperatures on Earth—is similar to today.

"Under a business-as-usual emissions scenarios, by 2100 warming in the upper [ocean](#) will have approached 20 percent of warming in the late Permian, and by the year 2300 it will reach between 35 and 50 percent," Penn said. "This study highlights the potential for a mass extinction arising from a similar mechanism under anthropogenic climate change."

More information: J.L. Penn et al., "Temperature-dependent hypoxia explains biogeography and severity of end-Permian marine mass extinction," *Science* (2018). [science.sciencemag.org/cgi/doi ... 1126/science.aat1327](https://science.sciencemag.org/cgi/doi/10.1126/science.aat1327)

Provided by University of Washington

Citation: Biggest mass extinction caused by global warming leaving ocean animals gasping for breath (2018, December 6) retrieved 9 April 2024 from <https://phys.org/news/2018-12-biggest-mass-extinction-global-ocean.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
