

Plant biologists identify mechanism behind transition from insect to wind pollination

December 17 2018



Representative examples of flowers with high (left) and low (right) natural frequency stamens. Flowers were stored in 60 percent ethanol and photographed using a dissection microscope. Credit: David Timerman

New research by scientists at the University of Toronto (U of T) offers novel insights into why and how wind-pollinated plants have evolved from insect-pollinated ancestors.

Early seed plants depended on wind to carry pollen between plants, but

about 100 million years ago, flowering plants evolved to attract insects that could transfer pollen with greater precision than random air currents. Although [insect pollination](#) is more economical, numerous lineages have since reverted back to wind pollination, leaving many biologists to question why that would ever happen given the success of insect pollination. This apparent paradox perplexed even Charles Darwin, and still today, little is known about the conditions initiating this transition.

In a study published this month in *Proceedings of the Royal Society B*, the researchers describe for the first time a mechanism driving this reversion involving the vibration of stamens, the pollen-bearing organs of flowers.

"We found that plants in which stamens vibrate more vigorously in wind, disperse pollen by wind more readily, and that this characteristic of stamens is favoured under conditions where plants receive few visits from pollinators," said lead author David Timerman, a Ph.D. candidate working with evolutionary biologist Spencer Barrett in the Department of Ecology & Evolutionary Biology in the Faculty of Arts & Science at U of T.

The discovery helps to explain the origins of wind pollination, which is represented in approximately 10 per cent of flowering [plant species](#).

"It may also be useful for understanding how plants will cope with a reduction in pollinator services due to the global collapse of wild pollinator populations," said Timerman.



Fly pollinator visiting a female flower of *Thalictum pubescens* (tall meadow-rue). Females of this species produce stamens with sterile pollen grains, presumably to reward pollinators. Credit: David Timerman

The reproductive structures of flowering plants are the most diverse of any group of living organisms. Flowers vary extensively in size, shape and structure, and much of this diversity is related to modes of pollination. Wind-pollinated species have independently evolved similar suites of floral traits adapted for releasing, dispersing and capturing pollen in air. One of these traits involves long flexible stamens that

vibrate conspicuously in wind relative to related insect-pollinated species.

Why and how wind pollination has evolved in flowering plants from animal pollination is a long-standing fundamental question in the evolutionary biology of plants. While wind pollination has evolved from animal pollination on at least 65 occasions in flowering plants—trees, ragweed and many grasses among them—the mechanisms involved in the transition are not well understood.

It has been long held by scientists that wind-pollinated plants are 'aerodynamically engineered' for efficient pollen dispersal. But compared to animal-pollinated species few studies have investigated the function of floral traits associated with wind pollination. Moreover, the modifications of flowers required for the evolutionary switch from insect to wind pollination have not been studied because until now, there have been no experiments on transitional species that are both wind and insect-pollinated

"We took a novel approach to this problem by applying biomechanics to understand the key processes involved in the early stages of this transition, and the work provided several novel insights," said Timerman.

Timerman and Barrett examined the problem in a species called *Thalictrum pubescens*, of the buttercup family. The plant is an ambophilous species, meaning it is pollinated by both insects and wind. As such, they speculate that the species probably represents a transitional state in the evolution of wind pollination.

Timerman used an electrodynamic shaker to apply controlled vibration to stamens to measure their natural frequency of vibration, then used a custom-built wind tunnel to investigate how the natural frequency of vibration influences pollen release. Timerman also performed a

manipulative field experiment at U of T's Koffler Scientific Reserve to confirm whether natural selection acts differently on stamen properties in the presence or absence of insect pollinators.



Fly pollinator approaching male flowers of *Thalictrum pubescens* (tall meadow-rue) to collect pollen. Credit: David Timerman

Timerman measured variation in the natural frequency of the stamen's vibration across nine populations, and assessed the repeatability of vibration frequency over consecutive growing seasons. With all of the data they collected, the researchers analysed the effect of this parameter on pollen release in the wind tunnel, as well as male reproductive success of plants in the field with and without pollinators.

"Successful reproduction was greatest for plants whose stamens vibrated

at a lower frequency when pollinators were absent, but this advantage diminished when pollinators were present," said Timerman. "Our biomechanical analysis of the wind-flower interface has identified this naturally occurring feature as a key trait for understanding early stages in the transition from insect to wind pollination."

Timerman says when animal pollinators do not provide adequate pollination services, natural selection should favour individuals with flexible stamens that vibrate readily, releasing pollen into the air.

"Wind is obviously a more consistent agent for pollen dispersal than relying on insects whose population sizes and behaviours fluctuate in time and space" says Timerman. "Further, many aspects of global environmental change are currently disrupting pollinator service to wild [plants](#), leading to what has been termed 'the pollination crisis'.

"These situations could potentially favour the evolution of [wind pollination](#) through the mechanism that we have discovered."

More information: *Proceedings of the Royal Society B* (2018). [DOI: 10.1098/rspb.2018.2251](https://doi.org/10.1098/rspb.2018.2251)

Provided by University of Toronto

Citation: Plant biologists identify mechanism behind transition from insect to wind pollination (2018, December 17) retrieved 18 April 2024 from <https://phys.org/news/2018-12-biologists-mechanism-transition-insect-pollination.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.