

Loss of intertidal ecosystem exposes coastal communities

December 19 2018



Sampling invertebrates within deep mudflats in Gladstone, Australia. Credit: Chi-Yeung Choi

Artificial intelligence and extensive satellite imagery have allowed



researchers to map the world's intertidal zones for the first time, revealing a significant loss of the crucial ecosystem.

The University of Queensland and University of New South Wales study has shown that global foreshore environments declined by up to 16 per cent between 1984 and 2016.

Professor Richard Fuller, from UQ's School of Biological Sciences, said the zone between low and high tide lines protected more than 625 million people around the world from storms and sea level rises.

"Identifying areas where intertidal zones are being lost to development and rising seas is critical to safeguard <u>coastal communities</u>," Professor Fuller said.

"Our research will have significant international benefits, with more than 1.4 billion people expected to live in <u>coastal areas</u> by 2060."

The study used <u>artificial intelligence</u> known as machine-learning to analyse more than 700,000 <u>satellite images</u> to map changing global distribution of intertidal areas over a 30-year period.

UQ and UNSW Faculty of Science researcher Dr. Nick Murray said it was made possible by a unique collaboration of conservation biologists, and marine, coastal, remote sensing, and computer scientists.





Dragging sediment samples across the mudflats in Gladstone, Australia. Credit: Chi-Yeung Choi

"It required nearly one million hours of computation, run on 22,000 machines via the Google Earth Engine," Dr. Murray said.

"We applied machine learning classifiers to every pixel from each satellite image available to us from along the world's coastlines."

The findings lay the ground-work for a global coastal monitoring system for international conservation and sustainable development targets.

"A system like this could enable scientists, governments and the wider community to take stock of the services that coastal ecosystems provide," Dr. Murray said.



"An online app, the Intertidal Change Explorer, offers <u>open access</u> to the dataset and supports its use to understand and conserve coastal ecosystems worldwide."



Intertidal loss in Incheon, South Korea. Credit: Nick Murray

The research, published in *Nature*, was supported by a philanthropic donation from Google.

More information: The global distribution and trajectory of tidal flats, *Nature* (2018). DOI: 10.1038/s41586-018-0805-8, www.nature.com/articles/s41586-018-0805-8



Provided by University of Queensland

Citation: Loss of intertidal ecosystem exposes coastal communities (2018, December 19) retrieved 27 June 2024 from <u>https://phys.org/news/2018-12-loss-intertidal-ecosystem-exposes-coastal.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.