


# Precision experiment first to isolate, measure weak force between protons, neutrons

December 19 2018, by Sara Shoemaker

---



Scientists analyzed the gamma rays emitted during the NPDGamma Experiment and found parity-violating asymmetry, which is a specific change in behavior in the force between a neutron and a proton. They measured a 30 parts per billion preference for gamma rays to be emitted antiparallel to the neutron spin when neutrons are captured by protons in liquid hydrogen. After observing that more gammas go down than up, the experiment resolved for the first time a mirror-asymmetric component or handedness of the weak force. Credit: Andy Sproles/Oak Ridge National Laboratory, U.S. Dept. of Energy

A team of scientists has for the first time measured the elusive weak interaction between protons and neutrons in the nucleus of an atom. They had chosen the simplest nucleus consisting of one neutron and one proton for the study.

Through a unique [neutron](#) experiment at the Department of Energy's Oak Ridge National Laboratory, experimental physicists resolved the [weak force](#) between the particles at the atom's core, predicted in the Standard Model that describes the elementary particles and their interactions.

Their result is sensitive to subtle aspects of the [strong force](#) between nuclear particles, which is still poorly understood.

The team's observation, described in *Physical Review Letters*, culminates decades of work performed with an apparatus known as NPDGamma. The first phase of the experiment took place at Los Alamos National Laboratory. Building on the knowledge gained at LANL, the team moved the project to ORNL to take advantage of the high neutron beam intensity produced at the lab's Spallation Neutron Source.

Protons and neutrons are made of smaller particles called quarks that are bound together by the strong interaction, which is one of the four known forces of nature: strong [force](#), electromagnetism, weak force and gravity. The weak force exists in the tiny distance within and between protons and neutrons; the [strong interaction](#) confines quarks in neutrons and protons.

The weak force also connects the axial spin and direction of motion of the nuclear particles, revealing subtle aspects of how quarks move inside protons and neutrons.

"The goal of the experiment was to isolate and measure one component of this weak interaction, which manifested as gamma rays that could be counted and verified with high statistical accuracy," said David Bowman, co-author and team leader for neutron physics at ORNL. "You have to detect a lot of gammas to see this tiny effect."

The NPDGamma Experiment, the first to be carried out at the Fundamental Neutron Physics Beamline at SNS, channeled cold neutrons toward a target of liquid hydrogen. The apparatus was designed to control the spin direction of the slow-moving neutrons, "flipping" them from spin-up to spin-down positions as desired. When the manipulated neutrons smashed into the target, they interacted with the protons within the liquid hydrogen's atoms, sending out gamma rays that were measured by special sensors.

After analyzing the gamma rays, the scientists found parity-violating asymmetry, which is a specific change in behavior in the force between a neutron and a [proton](#). "If parity were conserved, a nucleus spinning in the righthanded way and one spinning in the lefthanded way—as if they were mirrored images—would result in an equal number of gammas emitting up as emitting down," Bowman explained.

"But, in fact, we observed that more gammas go down than go up, which lead to successfully isolating and measuring a mirror-asymmetric component of the weak force."

The scientists ran the experiment numerous times for about two decades, counting and characterizing the [gamma rays](#) and collecting data from these events based on neutron spin direction and other factors.

The high intensity of the SNS, along with other improvements, allowed a count rate that is nearly 100 times higher compared with previous operation at the Los Alamos Neutron Science Center.

Results of the NPDGamma Experiment filled in a vital piece of information, yet there are still theories to be tested.

"There is a theory for the weak force between the quarks inside the proton and neutron, but the way that the strong force between the quarks translates into the force between the proton and the neutron is not fully understood," said W. Michael Snow, co-author and professor of experimental nuclear physics at Indiana University. "That's still an unsolved problem."

He compared the measurement of the weak force in relation with the strong force as a kind of tracer, similar to a tracer in biology that reveals a process of interest in a system without disturbing it.

"The weak interaction allows us to reveal some unique features of the dynamics of the quarks within the nucleus of an atom," Snow added.

**More information:** D. Blyth et al, First Observation of P -odd  $\gamma$  Asymmetry in Polarized Neutron Capture on Hydrogen, *Physical Review Letters* (2018). [DOI: 10.1103/PhysRevLett.121.242002](https://doi.org/10.1103/PhysRevLett.121.242002)

Provided by Oak Ridge National Laboratory

Citation: Precision experiment first to isolate, measure weak force between protons, neutrons (2018, December 19) retrieved 26 April 2024 from <https://phys.org/news/2018-12-precision-isolate-weak-protons-neutrons.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.