

Prescribed burning not as damaging as previously thought

December 3 2018



Prescribing burning at Moor House National Nature Reserve. Credit: University of Liverpool

New research by the University of Liverpool has found that prescribed burning, a controversial technique where fires are intentionally used to

manage vegetation, is not as damaging to peat growth as previously thought if carried out on a sensible rotation, and can produce several positive outcomes.

In a study published in *Nature Geoscience*, scientists analysed data from a long-term ecological experiment at Moor House National Nature Reserve which contained areas of moorland that only been burned in 1954, or since 1954 had burned either every ten or every twenty years. These were compared with "control" areas unburned since the 1920s.

Analysis of changes in vegetation composition, led by Emeritus Professor Rob Marrs from the University's Department of Earth, Oceans & Ecological Sciences, found that the areas which hadn't been subjected to any prescribed [burns](#) were dominated by heather, and other low-level, peat-forming [species](#), and contained less good peat-forming species such as Sphagnum mosses and cotton grasses.

New stratigraphical data collected on the rate of peat and carbon accumulation led by Professor Richard Chiverrell from the University's Department of Geography & Planning measured numerous peat profiles sampled from the different prescribed burn areas.

Professor Chiverrell said: " This is first time that stratigraphical techniques have been used within the structure of a designed experiment. Our data show only limited reduction of peat and carbon accumulation with increased burning treatments. Crucially, there was continued peat and carbon accumulation even in the areas that had undergone the regime of most frequent burning."

The researchers also modelled the effects of these experimental prescribed burning treatments on potential damage caused by severe wildfires, and predicted that prescribed burning on rotation in some areas would reduce fire fuel loads (the amount of flammable material

available to be burned in a fire), protect against more severe ecosystem damage, allow better access for fighting wildfires and promote faster recovery post-wildfire.

Prescribed burning is a technique used throughout the world however its use, especially on peatlands in Britain, is contentious. It has been thought by many involved in peatland conservation that prescribed burning stops peat growth, reduces carbon capture, increases coloration in water and reduces the amount of peat-forming species especially Sphagnum mosses and cotton grass.

Emeritus Professor Rob Marris said: "The research, which was truly interdisciplinary and combined several scientific approaches, centred on a long-term, replicated experiment at Moor House National Nature Reserve. As part of that experiment the moorland was subjected to four burning treatments to determine the change in composition of the main species. The value of long-term experiments is that they allow new hypotheses can be tested which were unforeseen at the start,

"Our study produced surprising results that are the opposite of what many conservationists had previously thought. It showed that the areas which had not been burnt since 1954 ended up with more heather and less peat forming species but the most frequently burned treatment had more of these species and did not stop carbon accumulation. We advise that prescribed burning be used in some areas within a moorland management plan to minimize fuel loads.

"This work is unique because it uses stratigraphical methods, usually used to measure change over millennia, to assess recent environmental changes within a relatively long-term manipulative field experiment."

More information: R. H. Marris et al, Experimental evidence for sustained carbon sequestration in fire-managed, peat moorlands, *Nature*

Geoscience (2018). [DOI: 10.1038/s41561-018-0266-6](https://doi.org/10.1038/s41561-018-0266-6)

Provided by University of Liverpool

Citation: Prescribed burning not as damaging as previously thought (2018, December 3) retrieved 24 April 2024 from <https://phys.org/news/2018-12-previously-thought.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.