


Proteins imaged in graphene liquid cell have higher radiation tolerance

December 10 2018



Schematic representation of hydrated microtubule proteins encapsulated between two graphene layers imaged by transmission electron microscopy (TEM). An example of a TEM image of a microtubule is shown at the right. The interior lining reflects the protofilament structure of the polymeric microtubule. Free within this press release. Credit: INM, Niels de Jonge

Electron microscopy is one of the main methods used to examine protein structure. Studying these structures is of key importance to elucidate their function feeding fundamental information into a number of fields such as structural biology, cell biology, cancer research, and other biomedical fields. It also enhances the understanding of biomineralization.

A new option for imaging proteins is liquid-phase [electron microscopy](#) (LPEM), which is capable of imaging native (unstained) [protein structure](#) and other samples such as nanomaterials or cells in liquid. This

technology was developed over the past fifteen years. Until recently, it debated whether the radiation tolerance of liquid samples would be better or worse compared to amorphous ice. In their recent publication, Sercan Keskin and Niels de Jonge from the INM-Leibniz Institute for New Materials now demonstrate, that the radiation tolerance is increased by an order of magnitude compared to a sample in ice. This result was achieved by preparing a microtubule sample in a graphene liquid cell. Essential was to use a low as possible rate at which the electron beam irradiation was applied.

Traditionally, samples were fixed, stained with a metal to enhance their contrast, subsequently dried, embedded in plastic, cut in thin sections, and then imaged in the vacuum environment required for electron microscopy. Cryo electron microscopy overcomes the drawbacks associated with this [sample](#) preparation and provides the means to study proteins in a close to native hydrated state by preparing them in amorphous ice. However, a key imitating is the high sensitivity of the samples to electron beam irradiation, so that statistical noise in the image prevents [high resolution](#) and many ten thousand noisy images of identical structures need to be imaged in order to resolve the structure.

More information: Sercan Keskin et al. Reduced Radiation Damage in Transmission Electron Microscopy of Proteins in Graphene Liquid Cells, *Nano Letters* (2018). [DOI: 10.1021/acs.nanolett.8b02490](https://doi.org/10.1021/acs.nanolett.8b02490)

Provided by Leibniz Institute for New Materials

Citation: Proteins imaged in graphene liquid cell have higher radiation tolerance (2018, December 10) retrieved 19 April 2024 from <https://phys.org/news/2018-12-proteins-imaged-graphene-liquid-cell.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.