


Scientists discover possible mantle mineral

December 7 2018



Maohokite. Credit: CHEN Ming

Scientists long believed that Earth's lower mantle was composed of Bridgmanite $(\text{Mg,Fe})\text{SiO}_3$ and magnesiowüstite $(\text{Mg,Fe})\text{O}$, in which Fe^{2+} dwells. This view changed when experiments showed that Fe^{2+} simply can't exist at the pressure and temperature of the lower mantle. What is present is Fe^{3+} . The two phases $(\text{Mg,Fe})\text{SiO}_3$ and $(\text{Mg,Fe})\text{O}$ both shed Fe^{2+} and, in turn, MgSiO_3 and MgO remain. However, what mineral hosts Fe^{3+} had remained unknown.

Now, scientists have a possible answer: Maohokite, a newly discovered high-pressure mineral. It may be what composes the Earth's lower mantle along with Bridgmanite MgSiO_3 and magnesiowüstite MgO . The study reporting this new mineral was published in *Meteoritics & Planetary Science*.

Maohokite was discovered by Chen Ming's team from the Guangzhou Institute of Geochemistry of the Chinese Academy of Sciences and SHU Jinfu from the Center for High Pressure Science and Technology Advanced Research. The mineral was named after Hokwang Mao, in honor of his great contribution to high-pressure research.

The mineral and its name have been approved by the Commission on New Minerals, Nomenclature and Classification of the International Mineralogical Association under the designator IMA 2017-047.

Natural minerals can be divided into two types: low-pressure minerals and high-pressure minerals, depending on their formation pressures. The pressure and temperature required for the formation of high-pressure minerals can only be provided by the environment of the mantle or the hypervelocity collision between celestial bodies.

Maohokite is the second case. It was found in shock-metamorphosed rocks from the Xiuyan impact crater in China.

This high-pressure mineral was formed from the decomposition of ferromagnesian carbonate via a self-oxidation-reduction reaction at a temperature >900 degrees C and impact pressure >25 GPa (a [pressure](#) range found at depths more than 670km below Earth's surface). In this reaction, Fe^{2+} oxidizes into Fe^{3+} and then later combines with Mg^{2+} to form maohokite, thus making it a possible important constituent of the lower mantle.

Maohokite, with a composition of MgFe_2O_4 , has an orthorhombic CaFe_2O_4 -type structure. The existing mineralogical model of the Earth's mantle shows that the ferromagnesian lower mantle is mainly composed of Bridgmanite $(\text{Mg,Fe})\text{SiO}_3$ and magnesiowüstite $(\text{Mg,Fe})\text{O}$. Therefore, the fact that Maohokite contains Mg and Fe, two major components of the lower mantle, only makes the case stronger that Maohokite is a key [mineral](#) in the lower [mantle](#).

More information: Ming Chen et al. Maohokite, a post-spinel polymorph of MgFe_2O_4 in shocked gneiss from the Xiuyan crater in China, *Meteoritics & Planetary Science* (2018). [DOI: 10.1111/maps.13222](#)

Provided by Chinese Academy of Sciences

Citation: Scientists discover possible mantle mineral (2018, December 7) retrieved 10 April 2024 from <https://phys.org/news/2018-12-scientists-mantle-mineral.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.