

# Scientists identify new minerals for carbon capture

December 12 2018, by Katie Willis

---



Sasha Wilson, associate professor in the Department of Earth and Atmospheric Sciences, supervised Turvey's research. The study identifies new minerals capable of carbon capture. Credit: John Ulan

Research confirms new minerals are capturing and storing carbon in a

new paper by University of Alberta geologists and their collaborators. The minerals, members of the hydrotalcite group, are the first outside of the carbonate family to naturally capture atmospheric CO<sub>2</sub> in mine waste, important as society continues to forge ways to lower our carbon emissions and combat climate change.

"This research confirmed that hydrotalcites are capable of sequestering atmospheric CO<sub>2</sub> in mine waste," said Connor Turvey, who conducted this research during his Ph.D. studies under the supervision of Sasha Wilson. "Hydrotalcites are trapping the CO<sub>2</sub> deeper into the tailings where carbonate minerals were unable to form."

Mine tailings are composed of the waste minerals removed from the ground in the mining process. As these minerals are exposed to the atmosphere and to rain water, they can react to form new minerals that trap CO<sub>2</sub> from the atmosphere.

The research highlights the potential for improving [carbon capture](#) one to two metres beneath the surface of mining wastes, where most sequestration usually occurs. "One thing that this indicates is that the capacity for [carbon sequestration](#) at this depth could be improved by providing a more plentiful source of [carbon](#) to those depths," added Wilson, associate professor in the Department of Earth and Atmospheric Sciences and expert in economic geology.

The study was conducted at Woodsreef Asbestos Mine in New South Wales, Australia. Carbon capture, utilization, and storage in minerals is of growing importance for both academia and industry. This discovery points to the potential to use mineral waste from mines to sequester carbon more effectively, supporting remediation efforts.

"Merely going [carbon neutral](#) is no longer going to be enough to prevent climate change from occurring," said Turvey. "What is now needed is

for our world to become carbon negative."

This means that industry must both reduce [carbon emissions](#) while simultaneously exploring carbon sequestration to actively draw excess CO<sub>2</sub> from the atmosphere—making discoveries like this one more important than ever.

The paper, "Hydrotalcites and hydrated Mg-carbonates as carbon sinks in serpentinite mineral wastes from the Woodsreef chrysotile mine, New South Wales, Australia: Controls on carbonate mineralogy and efficiency of CO<sub>2</sub> air capture in mine tailings," was published in the *International Journal of Greenhouse Gas Control*.

**More information:** Connor C. Turvey et al, Hydrotalcites and hydrated Mg-carbonates as carbon sinks in serpentinite mineral wastes from the Woodsreef chrysotile mine, New South Wales, Australia: Controls on carbonate mineralogy and efficiency of CO<sub>2</sub> air capture in mine tailings, *International Journal of Greenhouse Gas Control* (2018). [DOI: 10.1016/j.ijggc.2018.09.015](https://doi.org/10.1016/j.ijggc.2018.09.015)

Provided by University of Alberta

Citation: Scientists identify new minerals for carbon capture (2018, December 12) retrieved 11 May 2024 from <https://phys.org/news/2018-12-scientists-minerals-carbon-capture.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.