

Southwest forests may decline in productivity on average as much as 75 percent over the 21st century as climate warms, according to a University of Arizona-led research report published in *Nature Communications* on Dec. 17.

The new estimate is better than previous ones because it is based on a new database of information on the growth of trees under average conditions, according to the research team. Previous estimates were based on a database that included many trees growing in marginal conditions.

The finding is based on a treasure trove of about 20,000 unanalyzed tree cores discovered in a Utah laboratory about a decade ago. The annual growth rings visible in tree cores reflect each year's climatic conditions.

The new tree-[core](#) samples are more representative of the forest as a whole than many of those collected and analyzed earlier, said first author Stefan Klesse, who conducted the analysis while a [postdoctoral researcher](#) at the UA Laboratory of Tree-Ring Research.

Senior author Margaret Evans said the spatial representation in the new data set from the U.S. Forest Service Forest Inventory and Analysis program is "unprecedented." She calls the trees in the new data set that are living under average growing conditions "Joe Schmoe trees."

"The Joe Schmoe trees will experience a 75 percent reduction in growth and the trees on the edge—according to our analysis—are pretty much doomed," said Evans, an assistant professor in the UA Laboratory of Tree-Ring Research.

The database previously available to researchers is the International Tree-Ring Data Bank, or ITRDB, which included samples from many trees that were growing under marginal conditions—what Evans characterized

as "on the edge."



Tree cores mounted on sticks to make the cores easier to handle for measuring the annual growth rings. The numbers on the pink tags indicate in what county the cores were taken. Credit: John D. Shaw, US Forest Service

When the trees in the ITRDB were initially sampled, which was often decades ago, researchers chose trees living in marginal conditions because those trees were most sensitive to [climate](#) variation and thus best suited to reveal how climate varied over the past centuries.

Tree-ring researchers have long known the ITRDB samples might overestimate the effect of climate on average trees, but had no other data to use until now. And, even though the trees in the ITRDB may provide an overestimate of how the average forest tree will respond to the change in climate in the 21st century, that information is still valuable, Evans said.

"Those trees are the canaries in the coal mine—the edge of the forest is where we're going to see the change first," she said.


The research paper by Klesse, Evans and their co-authors, "Sampling bias overestimates climate change impacts on forest growth in the southwestern United States" was published in *Nature Communications* on Dec. 17. A list of co-authors is at the bottom of this release.

The U.S. Department of Agriculture, the U.S. Environmental Protection Agency and the Navajo Nation funded the research.

The new collection of Forest Inventory and Analysis tree cores turned up when scientists at the U.S. Forest Service Rocky Mountain Research

Station in Ogden, Utah, were moving from one building to another. The researchers were surprised to find a bunch of dusty boxes filled with tree cores that had never been analyzed.

The cores had been systematically collected from trees throughout the eight interior states of the U.S. West during the 1980s and 1990s. The cores reflected the growth of individual trees going back to the 1920s and some even earlier.



This map shows the geographic distribution of the tree-ring samples taken from forest trees in the US Four Corners states: Utah, Colorado, Arizona and New Mexico. Samples taken by the US Forest Service Forest Inventory and Analysis program are shown in blue. The samples taken and included in the International Tree-Ring Data Bank are shown in red. Credit: Stefan Klesse, Swiss Federal Research Institute WSL

Co-authors R. Justin DeRose and John Shaw of the U.S. Forest Service Rocky Mountain Research Station began the painstaking job of measuring [annual growth rings](#) from the approximately 20,000 cores, each from a specific tree. The two scientists enlisted Evans and sent her about 2,000 cores from Arizona trees. Her [research team](#), which included eight UA undergraduates, began cataloging those cores and recording the data from them in 2015.

The researchers wondered what the new tree-ring records might reveal about how climate change would affect the growth of forest trees of the Southwest in the latter half of the 21st century. To figure that out, they focused on records from common pinyon pine, Douglas fir and ponderosa pine from Utah, Colorado, Arizona and New Mexico.

Klesse, who is now a postdoctoral researcher at the Swiss Federal Research Institute WSL in Zurich, compared information from the newly analyzed cores from "Joe Schmoe" trees, plus additional tree-ring growth records from 858 trees in Arizona and New Mexico, with the records in the ITRDB.

For projections of the region's precipitation and temperature in the 21st century, he used climate projections from one of the most current

climate models, CMIP5 (Coupled Model Intercomparison Project Phase 5).

His analysis revealed that growth of the ordinary [forest](#) trees would not be reduced as much under climate change as the trees whose records are included in the ITRDB.

Even so, he said, "As the climate warms, tree [growth](#) will decline."

Evans said, "The [trees](#) have to take whatever they get in terms of climate conditions. When the temperatures rise they have to cope with it—or not."

More information: Stefan Klesse et al, Sampling bias overestimates climate change impacts on forest growth in the southwestern United States, *Nature Communications* (2018). [DOI: 10.1038/s41467-018-07800-y](#)

Provided by University of Arizona

Citation: Southwest forest trees will grow much slower in the 21st century (2018, December 18) retrieved 2 May 2024 from <https://phys.org/news/2018-12-southwest-forest-trees-slower-21st.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--