

Warming warning over turtle feminization

December 19 2018



Green turtle hatchling. Credit: University of Exeter

Up to 93% of green turtle hatchlings could be female by 2100, as climate change causes "feminisation" of the species, new research suggests.

The sex of turtle hatchlings is determined by [temperature](#), and at present

about 52% of hatching green [turtles](#)—one of seven species of sea turtle—are female.

But a study by the University of Exeter and the Marine and Environmental Sciences Centre (Portugal) shows that in [warmer temperatures](#) predicted by Intergovernmental Panel on Climate Change (IPCC) scenarios, 76-93% of hatchlings would be female.

The figures are specific to the study site in Guinea-Bissau, West Africa, but researchers say they expect a similar picture globally.

They say the changing gender ratio would initially lead to more females nesting, increasing the population, before by a decline "as incubation temperatures approach lethal levels".

They also predict rising sea levels will submerge 33-43% of current nesting areas used by green turtles on the beaches where the study was carried out.

"Green turtles are facing trouble in the future due to loss of habitats and increasing temperatures," said Dr. Rita Patricio, of the Centre for Ecology and Conservation on the University of Exeter's Penryn Campus in Cornwall.



Green turtle hatchling. Credit: University of Exeter

"Our results suggest the nesting population of green turtles the Bijagós Archipelago, Guinea-Bissau, will cope with the effects of [climate change](#) until 2100.

"Cooler temperatures, both at the end of the nesting season and in shaded areas, will guarantee some hatchlings are male.

"Although rising temperatures will lead to more female hatchlings—and 32-64% more nesting females by 2120—mortality in eggs will also be higher in these warmer conditions.

"As temperatures continue to rise, it may become impossible for

unhatched turtles to survive."

The [research team](#), which included the Institute of Biodiversity and Protected Areas of Guinea-Bissau, say nesting sites submerged by rising seas may not simply "move" inland.



Green turtle. Credit: University of Exeter

"Beach retreat may be OK in some areas, but the turtles we studied were nesting on a small island (Poilão Island) so there is a limit to how far the beach can go," Dr. Patricio said.

"In other places there may be natural barriers or human constructions that stop beaches moving inland."

The Bijagós Archipelago is the most important nesting place for [green turtles](#) in Africa, and the main breeding ground for the species in the South Atlantic.

The paper, published in the journal *Global Change Biology*, is entitled: "Climate change resilience of a globally important sea turtle nesting population."

Provided by University of Exeter

Citation: Warming warning over turtle feminization (2018, December 19) retrieved 19 April 2024 from <https://phys.org/news/2018-12-turtle-feminization.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.