


The algae's third eye

January 11 2019



In this multicellular Volvox alga, the novel light sensor 2c-cyclop was labeled

with fluorescence (green). It shows up in membranes around the nucleus. Credit: Eva Laura von der Heyde

Scientists at the Universities of Würzburg and Bielefeld in Germany have discovered an unusual new light sensor in green algae. The sensor triggers a reaction that is similar to one in the human eye.

Just like [land plants](#), algae use sunlight as an energy source. Many [green algae](#) actively move in the water; they can approach the [light](#) or move away from it. For this they use special sensors (photoreceptors) with which they perceive light.


The decades-long search for these light sensors led to a first success in 2002: Georg Nagel, at the time at Max-Planck-Institute of Biophysics in Frankfurt/M, and collaborators discovered and characterized two so-called channelrhodopsins in algae. These ion channels absorb light, then open up and transport ions. They were named after the visual pigments of humans and animals, the rhodopsins.

Now a third "eye" in algae is known: Researchers discovered a new light sensor with unexpected properties. The research groups of Professor Armin Hallmann (Bielefeld University) and Professor Georg Nagel (Julius-Maximilians-Universität Würzburg, JMU) report this finding in the journal *BMC Biology*.

Light reduces cGMP production

The surprise: The new photoreceptor is not activated by light but inhibited. It is a guanylyl cyclase which is an enzyme that synthesizes the important messenger cGMP. When exposed to light, cGMP production is severely reduced, leading to a reduced cGMP concentration – and

that's exactly what happens in the human eye as soon as the rhodopsins there absorb light.



Two molecules of the new light sensor 2c-cyclop, which in darkness produces

the messenger cGMP. Green light inhibits the signal chain. Credit: Shiqiang Gao / Armin Hallmann, Georg Nagel

The newly discovered sensor is regulated by light and by the molecule ATP. Such "two component systems" are already well known in bacteria, but not in higher evolved cells. The researchers have named the new photoreceptor "Two Component Cyclase Opsin", 2c-cyclop for short. They found it in two green algae, in the unicellular *Chlamydomonas reinhardtii* as well as in the multicellular *Volvox carteri*.

Function shown in oocytes and algae

"For many years there has been [genetic data](#) from which we could conclude that in green algae there must be many more rhodopsins than the two previously characterized," explains Georg Nagel. Only in *Chlamydomonas reinhardtii* twelve protein sequences are assigned to the opsins, which are the precursors of rhodopsins.

"So far, nobody could demonstrate the function of these light [sensors](#)," says Nagel's co-researcher Dr. Shiqiang Gao. Only the research groups from Bielefeld and Würzburg have succeeded in doing so: They have installed the new rhodopsin in oocytes of the toad *Xenopus laevis* and in the spherical alga *Volvox carteri*. In both cases, its function could be shown and characterized.

Perspectives for Optogenetics

The authors believe that the 2c-Cyclop light sensor offers new opportunities for optogenetics. With this methodology, the activity of living tissues and organisms can be influenced by light signals. By means of optogenetics, many basic biological processes in cells have already

been elucidated. For example, it provided new insights into the mechanisms of Parkinson's disease and other neurological diseases. She also brought new insights into diseases like autism, schizophrenia, and depression or anxiety disorders.

The JMU researcher Georg Nagel and the biophysicist Peter Hegemann (Humboldt Universität Berlin) are among the pioneers of optogenetics: They discovered the channelrhodopsins and found that the light-controlled [ion channels](#) from [algae](#) can be incorporated into animal cells and then controlled with light. For this achievement, both – together with other researchers – have received multiple awards.

More information: Yuehui Tian et al. Two-component cyclase opsins of green algae are ATP-dependent and light-inhibited guanylyl cyclases, *BMC Biology* (2018). [DOI: 10.1186/s12915-018-0613-5](https://doi.org/10.1186/s12915-018-0613-5)

Provided by Julius-Maximilians-Universität Würzburg

Citation: The algae's third eye (2019, January 11) retrieved 19 April 2024 from <https://phys.org/news/2019-01-algae-eye.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--