

# Why do beaked whales return to a Navy sonar range despite frequent disturbance?

January 29 2019

---



Cuvier's beaked whale on a Navy sonar range off California. Credit: A. Friedlaender; NMFS permit #14534

Using data from underwater robots, scientists have discovered that beaked whales prefer to feed within parts of a Navy sonar test range off Southern California that have dense patches of deep-sea squid. A new study published in the *Journal of Applied Ecology*, shows that beaked whales need these prey hotspots to survive, and that similar patches do not exist in nearby "sonar-free" areas.

For decades, the U.S. Navy has used high-powered sonar during anti-submarine training and testing exercises in various ocean habitats, including the San Nicolas Basin off Southern California. Beaked whales are particularly sensitive to these kinds of military sonars, which sometimes result in mass stranding events. Following [legal action](#) from environmental activists related to these risks, the Navy modified some training activities, created "sonar-free" areas, and spent more than a decade and tens of millions of dollars trying to find ways to reduce the harm to beaked whales and other mammals.

The new research, led by Brandon Southall at the University of California, Santa Cruz, and Kelly Benoit-Bird at the Monterey Bay Aquarium Research Institute, aimed to better understand why whales keep returning to the test range despite the risks.

The researchers equipped an underwater robot with echosounders to measure the abundance, density, and sizes of deep-sea squids in different parts of the Navy test range, as well as in nearby waters. They also developed an "energy budget" for beaked whales, showcasing the costs—in time and calories—of hunting for squid. This helped the researchers estimate how many dives the whales needed to make in order to get enough food to survive in different areas.

"Beaked whales work very hard to obtain their food. They are essentially living paycheck to paycheck," said Benoit-Bird. Unlike many [baleen whales](#) with significant energy reserves, beaked whales can't afford to expend too much energy on a dive that doesn't result in capturing many squid. In areas where the concentration of prey is low, the beaked whales must work harder and expend more calories—making reproduction and raising young that much more challenging. Some of the areas under study were so poor in terms of prey that whales likely could not meet their basic energetic requirements if they only fed there.

"Despite how things might look from the surface, the deep sea is not uniform," Benoit-Bird added. "There are pockets of wealth where squid are abundant, and beaked whales know exactly how to find those hot spots." It turns out that a portion of the Navy test range off Southern California encompasses one of these hot spots—an area rich in squid.

In fact, squid were 10 times more abundant higher in the area preferred by the whales. In this preferred area, the whales could get enough food by making just one dive a day. In a nearby sonar-free area (established with the idea that [beaked whales](#) could shelter in these areas while the sonar tests were underway) the [whales](#) would need to make between 22 and 100 dives per day to get enough food—something that would be difficult or impossible to do.

"Our findings, based on a novel integration of active sonar imaging and passive listening technologies, have multiple management implications," explained Southall. "They provide direct information to the Navy and federal regulators to better manage important and impacted habitat areas off California. And they give us new data on foraging ecology for population-level models of disturbance that have been at the heart of recent debate and litigation over spatial management and proposed sonar exclusion zones."

This study is the first to link habitat quality with beaked whale behavior in such fine spatial scales. It also demonstrates that scientists can't assess the quality of deep-sea habitats by simply making measurements at the ocean surface, or even by measuring the physical and chemical properties of the deep ocean. Direct measurements of the prey environment at the depths where animals are feeding, coupled with observations of when and where animals are foraging, are critical.

Until now, collecting such detailed data, even over small time and space scales, was virtually impossible. The researchers are now working on

tools that will help them study predators and prey over longer time periods, and in other areas where the Navy operates high-powered sonar. Similar field-research and modeling techniques could also be used to assess the potential impacts of other human activities that may disturb ocean animals, such as shipping traffic or offshore oil and gas development.

**More information:** Brandon L. Southall et al, Quantifying deep-sea predator-prey dynamics: Implications of biological heterogeneity for beaked whale conservation, *Journal of Applied Ecology* (2019). [DOI: 10.1111/1365-2664.13334](https://doi.org/10.1111/1365-2664.13334)

Provided by Monterey Bay Aquarium Research Institute

Citation: Why do beaked whales return to a Navy sonar range despite frequent disturbance? (2019, January 29) retrieved 25 April 2024 from <https://phys.org/news/2019-01-beaked-whales-navy-sonar-range.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---