


Climate change: Effect on sperm could hold key to species extinction

January 8 2019, by Kris Sales



Credit: CC0 Public Domain

Since the 1980s, [increasingly frequent and intense heatwaves](#) have contributed to more deaths than any other extreme weather event. The fingerprints of extreme events and climate change are widespread in the

natural world, where populations are showing stress responses.

A common fingerprint of a warmer world is a range shift, where the distribution of a [species](#) moves to higher altitudes or migrates toward the poles. A review of several hundred studies found an [average shift of 17km poleward, and 11 metres upslope](#), every decade. However, if [temperature changes](#) are too intense or lead species to geographic dead ends, local extinctions occur in the heat.

In 2003, [80 percent of relevant studies found the fingerprints](#) were seen among species, from grasses to trees and molluscs to mammals. Some migrated, some changed colour, some altered their bodies and some shifted their life cycle timings. A recent review of more than 100 studies found [8-50 percent of all species](#) will be threatened by [climate change](#) as a result.

High temperatures and extinctions

Currently, we have a [disturbingly limited knowledge](#) of which biological traits are sensitive to climate change and therefore responsible for local extinctions. However, a potential candidate is [male reproduction](#), because a range of medical and agricultural studies [in warm blooded animals](#) have shown that male infertility happens during heat stress.


However, until recently this had [rarely been explored outside fruit flies](#) in cold blooded animals. This is despite the fact that ectotherms – organisms that rely on heat in their environment to maintain a suitable body [temperature](#) – comprise most of biodiversity. Astonishingly, nearly [25 percent of all species](#) are thought to be a beetle.

The [red flour beetle](#) (*Tribolium castaneum*) is a useful ectotherm for [large experiments on reproduction](#), as they can go from egg to adult in a month at 30°C. Females can store male [sperm](#) in specialised organs

called spermathecae and they only need to keep 4 percent of a single ejaculate to enable them to produce offspring for up to 150 days.

To look at the [impact of heatwaves on reproduction](#), beetles were exposed to either standard control conditions or five-day heatwave temperatures, which were 5°C to 7°C above their preferred temperature. Afterwards, beetles mated and a variety of experiments looked for damage to their [reproductive success](#), sperm form and function, and offspring quality.

We found that 42°C heatwave temperatures halved the number of offspring males could produce relative to 30°C, with some males failing to produce any and mature sperm in female storage also experiencing damage from heatwaves. However, the reproductive output of pairs where only the females endured a five-day heatwave event was similar in all temperatures.



Heated males (B) had less sperm (stained green) and relatively more dead sperm (stained red) than control males (A). Credit: Kris Sales, Author provided

The decline was likely due to a combination of males becoming worse at mating, less sperm being transferred, less sperm transferred being alive, less sperm being kept in the females' spermathecae and more sperm being damaged and infertile.

Two results were particularly concerning. These beetles, and many cold-blooded animals, can live for years and are likely to see multiple heatwaves. When we exposed males to two [heatwave](#) events, ten days apart, their offspring production was less than 1 percent of that of unheated males.

This suggests that successive heatwaves can compound the damage of previous ones. The damage to offspring longevity and male fertility was another effect which was compounded over successive generations, and could lead to spiralling population declines.

Knowing what aspects of biology higher temperatures could compromise is essential to understanding how climate change affects nature. Hopefully, this new knowledge can help predict which species are most likely to be vulnerable, allowing conservationists to prepare for the trouble ahead.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Climate change: Effect on sperm could hold key to species extinction (2019, January 8) retrieved 18 April 2024 from <https://phys.org/news/2019-01-climate-effect-sperm-key-species.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.