


# Cosmic telescope zooms in on the beginning of time

January 9 2019



This is a Hubble Space Telescope image of a very distant quasar (at right) that has been brightened and split into three images by the effects of the gravitational field of a foreground galaxy (left). The crosses mark the centers of each quasar image. The quasar would have gone undetected if not for the power of gravitational lensing, which boosted its brightness by a factor of 50. The gravitational field of the foreground galaxy (seen at left) warps space like a

funhouse mirror, amplifying the quasar's light. Shining with the brilliance of 600 trillion suns, the quasar is fueled by a supermassive black hole at the heart of a young galaxy in the process of forming. The image shows the quasar as it looked 12.8 billion years ago - only about 1 billion years after the big bang. The quasar appears red because its blue light has been absorbed by diffuse gas in intergalactic space. By comparison, the foreground galaxy has bluer starlight. The quasar, cataloged as J043947.08+163415.7 (J0439+1634 for short), could hold the record of being the brightest in the early universe for some time, making it a unique object for follow-up studies. Credit: NASA, ESA, Xiaohui Fan (University of Arizona)

Observations from Gemini Observatory identify a key fingerprint of an extremely distant quasar, allowing astronomers to sample light emitted from the dawn of time. Astronomers happened upon this deep glimpse into space and time thanks to an unremarkable foreground galaxy acting as a gravitational lens, which magnified the quasar's ancient light. The Gemini observations provide critical pieces of the puzzle in confirming this object as the brightest appearing quasar so early in the history of the Universe, raising hopes that more sources like this will be found.

Before the cosmos reached its billionth birthday, some of the very first cosmic light began a long journey through the expanding Universe. One particular beam of light, from an energetic source called a quasar, serendipitously passed near an intervening galaxy, whose gravity bent and magnified the quasar's light and refocused it in our direction, allowing telescopes like Gemini North to probe the quasar in great detail.

"If it weren't for this makeshift cosmic [telescope](#), the quasar's light would appear about 50 times dimmer," said Xiaohui Fan of the University of Arizona who led the study. "This discovery demonstrates that strongly gravitationally lensed quasars do exist despite the fact that we've been looking for over 20 years and not found any others this far

back in time."


The Gemini observations provided key pieces of the puzzle by filling a critical hole in the data. The Gemini North telescope on Maunakea, Hawai'i, utilized the Gemini Near-InfraRed Spectrograph (GNIRS) to dissect a significant swath of the infrared part of the light's spectrum. The Gemini data contained the tell-tale signature of magnesium which is critical for determining how far back in time we are looking. The Gemini observations also led to a determination of the mass of the black hole powering the quasar. "When we combined the Gemini data with observations from multiple observatories on Maunakea, the Hubble Space Telescope, and other observatories around the world, we were able to paint a complete picture of the quasar and the intervening galaxy," said Feige Wang of the University of California, Santa Barbara, who is a member of the discovery team.

That picture reveals that the quasar is located extremely far back in time and space—shortly after what is known as the Epoch of Reionization—when the very first light emerged from the Big Bang. "This is one of the first sources to shine as the Universe emerged from the cosmic dark ages," said Jinyi Yang of the University of Arizona, another member of the discovery team. "Prior to this, no stars, quasars, or galaxies had been formed, until objects like this appeared like candles in the dark."

The foreground galaxy that enhances our view of the quasar is especially dim, which is extremely fortuitous. "If this galaxy were much brighter, we wouldn't have been able to differentiate it from the quasar," explained Fan, adding that this finding will change the way astronomers look for lensed quasars in the future and could significantly increase the number of lensed quasar discoveries. However, as Fan suggested, "We don't expect to find many quasars brighter than this one in the whole observable Universe."

The intense brilliance of the quasar, known as J0439+1634 (J0439+1634 for short), also suggests that it is fueled by a [supermassive black hole](#) at the heart of a young forming galaxy. The broad appearance of the magnesium fingerprint captured by Gemini also allowed astronomers to measure the mass of the quasar's supermassive black hole at 700 million times that of the Sun. The supermassive black hole is most likely surrounded by a sizable flattened disk of dust and gas. This torus of matter—known as an accretion disk—most likely continually spirals inward to feed the black hole powerhouse. Observations at submillimeter wavelengths with the James Clerk Maxwell Telescope on Maunakea suggest that the black hole is not only accreting gas but may be triggering star birth at a prodigious rate—which appears to be up to 10,000 stars per year; by comparison, our Milky Way Galaxy makes one star per year. However, because of the boosting effect of gravitational lensing, the actual rate of star formation could be much lower.

Quasars are extremely energetic sources powered by huge black holes thought to have resided in the very first [galaxies](#) to form in the Universe. Because of their brightness and distance, quasars provide a unique glimpse into the conditions in the early Universe. This quasar has a redshift of 6.51, which translates to a distance of 12.8 billion [light years](#), and appears to shine with a combined light of about 600 trillion Suns, boosted by the gravitational lensing magnification. The foreground galaxy which bent the quasar's light is about half that distance away, at a mere 6 billion light years from us.



The light from quasar J0439+1634, some 12.8 billion light years away, passes close to a faint galaxy that is about six billion light years away. The gravity of this foreground galaxy warps the space around it, according to Einstein's theory of general relativity. This bends the light like an optical lens, magnifies the quasar image by a factor of fifty, while at the same time split the quasar image into three. Both the foreground galaxy and the multiply imaged quasar is captured by the high resolution image of the Hubble Space Telescope. Ground-based telescopes, including the MMT, Keck, Gemini, LBT and JCMT, are used to observe this object in optical, infrared and sub-millimeter wavelengths to measure its distance, and to characterize its central black hole and host galaxy. Credit: NASA, ESA, Xiaohui Fan (University of Arizona)

Fan's team selected J0439+1634 as a very distant quasar candidate based

on optical data from several sources: the Panoramic Survey Telescope and Rapid Response System1 (Pan-STARRS1; operated by the University of Hawai'i's Institute for Astronomy), the United Kingdom Infra-Red Telescope Hemisphere Survey (conducted on Maunakea, Hawai'i), and NASA's Wide-field Infrared Survey Explorer (WISE) space telescope archive.

The first follow-up spectroscopic observations, conducted at the Multi-Mirror Telescope in Arizona, confirmed the object as a high-redshift quasar. Subsequent observations with the Gemini North and Keck I telescopes in Hawai'i confirmed the MMT's finding, and led to Gemini's detection of the crucial magnesium fingerprint—the key to nailing down the quasar's fantastic distance. However, the foreground lensing galaxy and the quasar appear so close that it is impossible to separate them with images taken from the ground due to blurring of the Earth's atmosphere. It took the exquisitely sharp images by the Hubble Space Telescope to reveal that the quasar image is split into three components by a faint lensing galaxy.

The quasar is ripe for future scrutiny. Astronomers also plan to use the Atacama Large Millimeter/submillimeter Array, and eventually NASA's James Webb Space Telescope, to look within 150 light-years of the black hole and directly detect the influence of the gravity from black hole on gas motion and star formation in its vicinity. Any future discoveries of very distant quasars like J0439+1634 will continue to teach astronomers about the chemical environment and the growth of massive black holes in our early Universe.

The study is described in a presentation at the 233rd meeting of the American Astronomical Society in Seattle, Washington and published in *The Astrophysical Journal Letters*.

**More information:** Xiaohui Fan et al. The Discovery of a

Gravitationally Lensed Quasar at  $z = 6.51$ , *The Astrophysical Journal* (2019). DOI: 10.3847/2041-8213/aaeffe

Provided by Gemini Observatory

Citation: Cosmic telescope zooms in on the beginning of time (2019, January 9) retrieved 25 April 2024 from <https://phys.org/news/2019-01-cosmic-telescope.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--