


Stick insect study shows the significance of passive muscle force for fast movements

January 10 2019



While the central nervous system has to actively control fast movements of large limbs, passive muscle force can suffice for the movement velocity and movement amplitude of small and light limbs. Credit: Arndt von Twickel *et al.*

Long, heavy limbs such as arms or legs differ fundamentally from short, light limbs such as fingers in their ability to execute fast movements. While the central nervous system has actively controls fast movements of large limbs, passive muscle force can suffice for the movement velocity and movement amplitude of small and light limbs. That is the result of a study conducted on the stick insect by the zoologists Ansgar Bueschges, Arndt von Twickel and Christoph Guschlbauer at the University of Cologne in cooperation with the visiting scientist Scott Hooper from Ohio University. The paper, "Swing Velocity Profiles of

Small Limbs Can Arise from Transient Passive Torques of the Antagonist Muscle Alone," has been published in *Current Biology*.

There are different basic speed levels in the course of a movement until the entire motor action has been optimally executed, says Bueschges. "If we look at the swinging of our legs when we walk, for example, the swinging phase brings the leg back into the starting position for the next step. With this feedback, the swing speed of the leg decreases toward the end, so that the foot does not touch down as quickly. Without the deceleration, the force generated when touching down would be so strong that it would counteract the next step." The [nervous system](#) produces this deceleration in large limbs such as the human leg.

For smaller limbs such as insect legs, however, the authors have shown that the slowing of the leg during rapid movements such as the swinging phase is independent of the [central nervous system](#). Von Twickel explains that this is due to the intrinsic muscle characteristics of the [limb](#) : "If the extensor muscle of a leg joint is actively shortened to produce rapid movements, the inactive flexor muscle is necessarily lengthened. During this stretching, the flexor muscle develops a previously unknown dynamic passive force, which is so large that it can continuously slow down movements to the functionally necessary level."

In the study, the authors exposed the flexor muscle of the [stick insect](#) leg to various stretching scenarios and measured the dynamic forces exerted by the passive muscle against the different stretches. Then, they used the results to dynamically simulate the functioning of a leg joint. The passive dynamic forces of the flexor generated by extensor activation were able to generate a movement that corresponded to the speed profile of an animal.

These results are quite surprising and contradict widespread notions of how small limbs execute fast movements. Bueschges concludes: "We

have known for a long time that passive muscle forces are important for the movements of small limbs, but we did not expect that their effect is so great that they can determine the speed profile of a [movement](#). This means that the active [muscle](#) and its opponent must be perfectly tuned to each other. We are in the process of better understanding this interaction."

More information: Arndt von Twickel et al, Swing Velocity Profiles of Small Limbs Can Arise from Transient Passive Torques of the Antagonist Muscle Alone, *Current Biology* (2018). [DOI: 10.1016/j.cub.2018.11.016](#)

Provided by University of Cologne

Citation: Stick insect study shows the significance of passive muscle force for fast movements (2019, January 10) retrieved 1 May 2024 from <https://phys.org/news/2019-01-insect-significance-passive-muscle-fast.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
