

Researchers observe inwardly rotating spirals in a nonoscillatory medium

January 29 2019, by Ingrid Fadelli



Credit: H. N. Yoshikawa



A team of researchers at Université Côte d'Azur and Hokkaido University have recently carried out a study exploring the spontaneous formation of spiral patterns observed on the downward-facing free surface of a horizontal liquid film. The surface examined by them entails Rayleigh-Taylor instability, which destabilizes the interface between two fluids of different densities when the heavier fluid is pushing down the lighter one.

The researchers observed that the liquid discharge resulting from this instability can occur in the form of propagating liquid curtains, which are generated at the film's circular periphery and appear as inwardly rotating spiral arms. Using a phenomenologically constructed cellular automaton, they demonstrated that these patterns arise from the phase locking, which leads to an intermittent discharge of liquid at a constant flow rate over the entire film surface.

"About 15 years ago, Dr. Laurent Limat and his colleagues studied spatiotemporal dynamics of liquid columns moving freely along the edge of a circular dish," Christian Mathis, one of the researchers who carried out the present study, told *Phys.org.* "Inspired by their work, we began the study of a bi-dimensional assembly of liquid columns, hoping to describe all the secondary instabilities related to losses of symmetry. We just found one such instability, but we discovered that we were in front of a rather simple system showing extremely rich behavior."

In their study, Mathis and his colleagues observed that increasing the flow rate gave rise to a series of complex patterns, including a regular hexagonal lattice of drops, a regular hexagonal lattice of columns, spatiotemporally intermittent behavior of columns and, finally, liquid curtains forming spiral waves.



The apparatus they used is fairly simple, consisting primarily of a semiclosed cylindrical vessel, with silicone oil continuously pouring into it through an inlet at the top and a fine mesh grid at the bottom. The vessel contains a constant amount of oil, determined by the equilibrium of a negative pressure in the vessel and the weight of the oil.

Excessive liquid in the vessel leaks through the grid and forms a liquid film underneath it. This film destabilizes in different ways, depending on the oil flow-rate and viscosity. Inside this apparatus, everything is made of transparent PMMA, as this enables easy observation and precise video measurements with adapted lightning.



Credit: H. N. Yoshikawa



"At a sufficiently high <u>flow-rate</u>, liquid curtains appear, move like waves and, end up coiling around some center point and forming rotating spiral arms," Mathis explained. "We measured the frequency of rotation of each arm and of the whole pattern and the length of arms. The key observation is the spontaneous formation of spirals through breaking the rotational symmetry of the system."

Spiral patterns can be found in many systems, both in lab experiments and in nature. Most of these spirals patterns rotate, with the <u>spiral arms</u> trailing the direction of this rotation. Patterns that rotate in the opposite direction are quite rare and are referred to as antispirals.

"We have shown that antispiral patterns can be produced in a simple liquid system, for which we know all the governing equations," Harunori Yoshikawa, another researcher who carried out the study, told *Phys.org*. "Deep investigation of this 'well-known' system would provide insights into rare occurrences of antispiral patterns."

The experiment carried out by Yoshikawa, Mathis and their colleagues could pave the way for further studies exploring the manifestation of antispiral patterns. In addition, their study could provide a benchmark for theories of pattern formation.

"We are now focusing on theoretical modeling, seeking an appropriate model equation and the ranges of involved parameters," Yoshikawa said. "We hope that we will theoretically reveal the essential features of these pattern formation processes."

The researchers are also currently exploring other patterns observed in their system, as the inwardly rotating spirals were merely one of their observations. For instance, the dripping regime and some of the peculiar behavior of the liquid columns are yet to be extensively investigated. In the study carried out by Mathis, Yoshikawa and their colleagues, the



dynamic of the spirals appeared to be related to the characteristics of the liquid film itself, a finding that they are also looking to explore further.

More information: Harunori N. Yoshikawa et al. Inwardly Rotating Spirals in a Nonoscillatory Medium, *Physical Review Letters* (2019). DOI: 10.1103/PhysRevLett.122.014502

© 2019 Science X Network

Citation: Researchers observe inwardly rotating spirals in a nonoscillatory medium (2019, January 29) retrieved 22 May 2024 from <u>https://phys.org/news/2019-01-inwardly-rotating-spirals-nonoscillatory-medium.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.