


Membraneless protocells could provide clues to formation of early life

January 31 2019, by Sam Sholtis



Membraneless protocells -- called complex coacervates -- can bring together molecules of RNA allowing the RNAs to perform certain reactions, an important step in the origin of life on Earth. The Image shows droplets of complex coacervates as seen under a microscope. The inset shows RNA molecules (cyan) are highly concentrated inside the droplets compared to the surrounding (dark). At roughly 2-5 micrometers in diameter, the droplets are about 14-35 times thinner than human hair. Credit: Bevilacqua Laboratory, Penn State

Membraneless assemblies of positively- and negatively-charged molecules can bring together RNA molecules in dense liquid droplets, allowing the RNAs to participate in fundamental chemical reactions. These assemblies, called "complex coacervates," also enhance the ability of some RNA molecules themselves to act as enzymes—molecules that drive chemical reactions. They do this by concentrating the RNA enzymes, their substrates, and other molecules required for the reaction. The results of testing and observation of these coacervates provide clues to reconstructing some of the early steps required for the origin of life on Earth in what is referred to as the prebiotic "RNA world." A paper describing the research, by scientists at Penn State, appears January 30, 2019 in the journal *Nature Communications*.

"We're interested in how you go from a world with no life to one with life," said Philip C. Bevilacqua, Distinguished Professor of Chemistry and of Biochemistry and Molecular Biology at Penn State and one of the senior authors of the paper. "One can imagine a lot of steps in this process, but we are not looking at the most elemental steps. We are interested in a slightly later step, to see how RNA [molecules](#) could form from their basic building blocks and if those RNA molecules could drive the reactions needed for life in the absence of proteins."

Life as we know it today generally requires genetic material—DNA, which is first transcribed into RNA. These two molecules carry information for the production of proteins, which are in turn required for most functional aspects of life, including the production of new [genetic material](#). This sets up a "chicken and the egg" dilemma for the origins of life on early Earth. DNA is required to produce proteins, but proteins are required to produce DNA.

"RNA—or something similar—has been thought of as a key to solving

this dilemma," said Raghav R. Poudyal, Simons Origins of Life Postdoctoral Fellow at Penn State and first author of the paper. "RNA molecules carry genetic information, but they can also function as enzymes to catalyze the [chemical](#) reactions needed for early life. This fact has led to the notion that life on Earth went through a stage where RNA played an active role in facilitating chemical reactions—"the RNA World"—where self-replicating RNA molecules both carried the genetic information and performed functions that are now generally carried out by proteins."

Another common feature of life on Earth is that it is compartmentalized in cells, often with an outer membrane, or in smaller compartments inside cells. These compartments ensure that all the components for the chemical reactions of life are in easy reach, but in the prebiotic world the building blocks for RNA—or the RNA enzymes needed to drive the chemical reactions that could lead to life—would probably have been scarce, floating around in the primordial soup.

"You can think of these RNA enzymes like a car being produced in an assembly line," said Poudyal. "If you don't have the parts in the right place in the factory, the assembly line doesn't work. Without coacervates, the parts needed for chemical reactions are too dilute and are unlikely to find each other, but inside the coacervates, all the parts that the enzyme needs to work are nearby."

The researchers therefore looked at a variety of materials that may have existed in the pre-life Earth that can form coacervates—membraneless protocells—and then allowed critical functions like sequestering the building blocks of RNA and bringing together RNA enzymes and their targets.

"It was previously known that RNA molecules can assemble and elongate in solutions with high concentrations of magnesium," said

Poudyal. "Our work shows that coacervates made from certain materials allow this non-enzymatic template-mediated RNA assembly to occur even in the absence of magnesium."

The coacervates are composed of positively charged molecules called polyamines and negatively charged polymers which cluster together to form membraneless compartments in a solution. Negatively charged RNA molecules are also attracted to the polyamines in the coacervates. Within the coacervates the RNA molecules are as much as 4000 times more concentrated than in the surrounding solution. By concentrating the RNA molecules in the coacervates, RNA enzymes are more likely to find their targets to drive chemical reactions.

"Although all the polyamines we tested were able to participate in formation of RNA-rich droplets, they differed in their ability to support RNA elongation," said Christine Keating, professor of chemistry at Penn State and a senior author on the paper. "These observations help us understand how the chemical environment within different membraneless compartments can impact RNA reactions."

"Although we can't look back to see the exact steps taken to form the first life on Earth, coacervates like the ones we can create in the laboratory may have helped by facilitating [chemical reactions](#) that otherwise would not have been possible," said Poudyal.

In addition to Bevilacqua, Poudyal, and Keating, the research team at Penn State includes Rebecca M. Guth-Metzler, Andrew J. Veenis, and Erica A. Frankel. The research was supported by the Simons Foundation and NASA.

More information: Raghav R. Poudyal et al, Template-directed RNA polymerization and enhanced ribozyme catalysis inside membraneless compartments formed by coacervates, *Nature Communications* (2019).

[DOI: 10.1038/s41467-019-08353-4](https://doi.org/10.1038/s41467-019-08353-4)

Provided by Pennsylvania State University

Citation: Membraneless protocells could provide clues to formation of early life (2019, January 31) retrieved 28 April 2024 from <https://phys.org/news/2019-01-membraneless-protocells-clues-formation-early.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.