

# Missing-link in planet evolution found

January 28 2019

---



Artist's impression of the newly discovered object. Credit: Ko Arimatsu

For the first time ever, astronomers have detected a 1.3 km radius body at the edge of the solar system. Kilometer-sized bodies like the one discovered have been predicted to exist for more than 70 years. These

objects acted as an important step in the planet formation process between small initial amalgamations of dust and ice and the planets we see today.

The Edgeworth-Kuiper Belt is a collection of small celestial bodies located beyond Neptune's orbit. The most famous Edgeworth-Kuiper Belt Object is Pluto. Edgeworth-Kuiper Belt Objects are believed to be remnants left over from the formation of the solar system. While small bodies like asteroids in the inner solar system have been altered by solar radiation, collisions, and the gravity of the planets over time; objects in the cold, dark, lonely Edgeworth-Kuiper Belt preserve the pristine conditions of the early solar system. Thus astronomers study them to learn about the beginning of the planet formation process.

Edgeworth-Kuiper Belt objects with radii from 1 kilometer to several kilometers have been predicted to exist, but they are too distant, small, and dim for even world-leading telescopes, like the Subaru Telescope, to observe directly. So a research team led by Ko Arimatsu at the National Astronomical Observatory of Japan used a technique known as occultation: monitoring a large number of stars and watching for the shadow of an object passing in front of one of them. The Organized Autotelescopes for Serendipitous Event Survey (OASES) team placed two small 28 cm telescopes on the roof of the Miyako open-air school on Miyako Island, Miyakojima-shi, Okinawa Prefecture, Japan, and monitored approximately 2000 stars for a total of 60 hours.

Analyzing the data, the team found an event consistent with a star appearing to dim as it is occulted by a 1.3 km radius Edgeworth-Kuiper Belt Object. This detection indicates that kilometer sized Edgeworth-Kuiper Belt Objects are more numerous than previously thought. This supports models where planetesimals first grow slowly into kilometer sized objects before runaway growth causes them to merge into [planets](#).

Arimatsu explains, "This is a real victory for little projects. Our team had less than 0.3 percent of the budget of large international projects. We didn't even have enough money to build a second dome to protect our second telescope! Yet we still managed to make a discovery that is impossible for the big projects. Now that we know our system works, we will investigate the Edgeworth-Kuiper Belt in more detail. We also have our sights set on the still undiscovered Oort Cloud out beyond that."

The study is published in *Nature Astronomy*.

**More information:** A kilometre-sized Kuiper belt object discovered by stellar occultation using amateur telescopes, *Nature Astronomy* (2019). [DOI: 10.1038/s41550-018-0685-8](https://doi.org/10.1038/s41550-018-0685-8) , [www.nature.com/articles/s41550-018-0685-8](https://www.nature.com/articles/s41550-018-0685-8)

Provided by National Institutes of Natural Sciences

Citation: Missing-link in planet evolution found (2019, January 28) retrieved 25 April 2024 from <https://phys.org/news/2019-01-missing-link-planet-evolution.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--