

Orchards in natural habitats draw bee diversity, improve apple production

January 17 2019



Wild bee in the genus *Melandrena* collecting pollen from an apple flower.
Credit: Kent Loeffler

Apple orchards surrounded by agricultural lands are visited by a less diverse collection of bee species than orchards surrounded by natural

habitats, according to a new Cornell University-led study, published in the journal *Science*.

In turn, apple production suffers when fewer, more closely-related species of bees pollinate an orchard. Production improves in orchards surrounded by [natural habitats](#), which then draw a broader selection of species to apple blossoms.

The researchers examined 10 years of data from 27 New York state [apple orchards](#). The study accounted for the types of landscapes that surround these orchards, measured apple production and surveyed the species of bees that visited each orchard.

The researchers also reconstructed the [evolutionary history](#) and relatedness of New York native bee species to better understand species patterns that played out across these orchard bee communities. This reconstruction is represented by a branching tree-like diagram of related species, called a phylogeny.

"Orchards that have bee communities that are more closely related to each other did worse in terms of their fruit production, and the communities that are more broad across the phylogeny did much better," said Heather Grab, Ph.D., the paper's first author and a postdoctoral researcher in the lab of Katja Poveda, associate professor of entomology and a co-author of the study. Brian Danforth, professor of entomology, is a senior author of the study.

Species of bees exhibit different behaviors in how and when they pollinate flowers. Some [species](#) approach from the side, others from the top, and they each may feed at different times of day and with varied frequencies, all of which affect how completely an apple flower is pollinated.

Organs in apple flowers must receive a certain number of pollen grains in order to develop a full complement of seeds. When seeds do well, the tissue that supports those seeds, the fleshy part of the fruit, is also more fully developed.

"If only half of the seeds mature fully, then the fruit is misshapen," which in turn affects weight and salability, Grab said.

In this way, habitats that surround farms affect the diversity of bee communities and, thus, an [orchard](#)'s productivity.

More information: H. Grab et al., "Agriculturally dominated landscapes reduce bee phylogenetic diversity and pollination services," *Science* (2019). [science.sciencemag.org/cgi/doi ... 1126/science.aat6016](https://science.sciencemag.org/cgi/doi/10.1126/science.aat6016)

Provided by Cornell University

Citation: Orchards in natural habitats draw bee diversity, improve apple production (2019, January 17) retrieved 6 February 2023 from <https://phys.org/news/2019-01-orchards-natural-habitats-bee-diversity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.