


New study quantifies deep reaction behind 'superdeep' diamonds

January 17 2019, by Joshua Rapp Learn





The Cullinan Diamond, the largest gem-quality diamond found, was discovered in South Africa in 1905. Superdeep diamonds have been uncovered at the same mine. Credit: Public Domain



Whether they are found in an engagement ring or an antique necklace, diamonds usually generate quick reactions from their recipients. Now, new research shows deep inside the Earth, fast reactions between subducted tectonic plates and the mantle at specific depths may be responsible for generating the most valuable diamonds.

The diamonds mined most often around the world are formed in the Earth's mantle at depths of around 150-250 kilometers (93-155 miles). They are created by <u>extreme pressure</u> and temperature of at least 1050 degrees Celsius (1922 degrees Fahrenheit). Only a small amount of these diamonds make it to mineable regions since most are destroyed in the process of reaching the Earth's crust via deep source volcanic eruptions.

But a tiny portion of mined diamonds, called sub-lithospheric or superdeep diamonds, are formed at much deeper depths than others, mostly in two rich zones at depths of 250-450 kilometers (155-279 miles) and 600-800 kilometers (372-497 miles). These diamonds stand out from others due to their compositions, which occasionally include materials from the deep Earth like majorite garnet, ferropericlase and bridgmanite.

"Although only composing 1 percent of the total mined diamonds, it seems lots of large and high-purity diamonds are superdeep diamonds, so they have good value as gems," said Feng Zhu the lead author of the new study in *Geophysical Research Letters*, a journal of the American Geophysical Union, who was a post-doctoral geology researcher at the University of Michigan when he performed the research.

No previous theory has completely explained the reason why very few diamonds have been found near the surface from the area at depths of 450-600 kilometers (372-497 miles) – the <u>region</u> between the zones where most superdeep diamonds are formed.



The new study seeks to explain this phenomenon. Zhu, now a post-doctoral researcher at the University of Hawai'i, and his colleagues believe the two superdeep areas where diamonds are formed are rich in the gems due to high production rates. The new study explains what drives the diamond-producing reaction in some areas and what slows it down in other areas.

Diamond formation

According to the authors, diamonds can form anywhere in the mantle, which extends from about 35 to 2,890 kilometers (21-1,800 miles) below the Earth's surface. However, humans rarely see most of the diamonds formed. Very few diamonds survive the volcanic trip to the Earth's crust where we can sample them.

That means the chances of finding diamonds from deep regions in the mantle, which produce relatively few of the gems, is extremely small. Only 1 percent of mined diamonds come from superdeep regions.

"In our hypothesis, the production of diamonds at any depth in the mantle is possible, it's just the production rate is different, so they have a different chance to be sampled in the crust," Zhu said.

Creating diamonds

In order to mimic the extreme pressures experienced deep inside planets, the study's authors used diamond anvil cells and a 1,000-ton multi-anvil apparatus at the University of Michigan. Both these devices allow researchers to compress sub-millimeter-sized material in extreme pressures. They compressed magnesium carbonate powder with iron foil in extreme heats and managed to create minuscule diamond grains visible through scanning electron microscopes.



They found that when conditions are right, diamond grains can form as quickly as every couple of minutes, and never took longer than a few hours to form, although the growth of gem diamonds may take much longer time in an actual melting fluid environment.

In the shallower region rich in superdeep diamond formation, 250-450 kilometers (155-279 miles) down, a subducting tectonic plate pushes under the Earth's mantle. This supplies plenty of carbonate, which creates "factories on a conveyor belt" for diamonds when combined with the iron from the mantle, the authors said.

High temperatures promote reactions which form diamonds, but pressure does the opposite. At depths roughly 475 kilometers (295 miles) below the surface, the pressure increases, and the reactions slow down drastically, the authors said. That's why few <u>diamonds</u> are found near the Earth surface coming from between 450-600 kilometers (372-497 miles).

"When your pressure reaches the diamond stable region, it will form. But when you increase pressure it will form at lower rates. You have a trade off there," Zhu said.

One exception to this rule is in the deeper region of 600-800 kilometers (372-497 miles) beneath the surface. In this region, accumulation of carbonate due to the stagnation of tectonic slabs pushing downwards makes up for the overdose in pressure. So while the reactions slow down, higher temperatures and an abundance of carbonate makes for a diamond-rich region.

Zhu said the new study adds to scientists' understanding of the Earth's mantle, about which relatively little is known for sure.

"Superdeep diamond inclusions bring us the only mineral samples from



the Earth's deep mantle," he said. "Seeing is believing, and these inclusions provide a solid ground for the studies on the inaccessible mantle."

More information: Feng Zhu et al. Kinetic control on the depth distribution of superdeep diamonds, *Geophysical Research Letters* (2018). DOI: 10.1029/2018GL080740

This story is republished courtesy of AGU Blogs (http://blogs.agu.org), a community of Earth and space science blogs, hosted by the American Geophysical Union. Read the original story here.

Provided by American Geophysical Union

Citation: New study quantifies deep reaction behind 'superdeep' diamonds (2019, January 17) retrieved 20 March 2024 from https://phys.org/news/2019-01-quantifies-deep-reaction-superdeep-diamonds.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.