


Second scientific balloon launches from Antarctica

January 2 2019, by Talia Ogliore



Panels are loaded onto X-Calibur in preparation for launch from McMurdo Station, Antarctica. Credit: Dana Braun

Washington University in St. Louis announced that its X-Calibur

instrument, a telescope that measures the polarization of X-rays arriving from distant neutron stars, black holes and other exotic celestial bodies, launched today from McMurdo Station, Antarctica.

The telescope is carried aloft on a helium balloon intended to reach an altitude of 130,000 feet. At this height, X-Calibur will travel at nearly four times the cruising altitude of commercial airliners, and above 99 percent of the Earth's atmosphere.

"Our prime observation target will be Vela X-1, a neutron star in binary orbit with a supergiant star," said Henric Krawczynski, professor of physics in Arts & Sciences at Washington University. The team hopes to gain new insights into how [neutron stars](#) and [black holes](#) in a binary orbit with stars grow by gobbling up stellar matter.

Researchers will combine observations from the balloon-borne X-Calibur with simultaneous measurements from three existing, space-based satellites.

"The results from these different observatories will be combined to constrain the [physical conditions](#) close to the neutron star, and thus to use Vela X-1 as a laboratory to test the behavior of matter and magnetic fields in truly extreme conditions," Krawczynski said.

X-Calibur will need to spend at least eight days aloft to gather enough data for scientists to consider it a success. During this time, the balloon is expected to make a single revolution around the Antarctic continent. If conditions permit, X-Calibur may be flown for additional days.



The harsh environment around McMurdo Station, Antarctica, prior to attempts to launch X-Calibur by a team of Washington University researchers. Credit: Fabian Kislat

X-Calibur is designed to measure the polarization—or, roughly, the orientation of the electric field—of incoming X-rays from binary systems.

Researchers hope to use the Vela X-1 observations to reveal how neutron stars accelerate particles to high energies. The observations furthermore will test two of the most important theories in modern physics under [extreme conditions: quantum electrodynamics](#) and general relativity.

Quantum electrodynamics predicts that the quantum vacuum close to

magnetized neutron stars exhibits birefringent properties—that is, it affects X-rays in a similar way as birefringent crystals such as sapphires or quartz affect optical light.

The theory of general relativity describes the trajectories of the X-rays close to the neutron stars where the extreme mass of the [neutron](#) stars almost curves spacetime into a knot.

Provided by Washington University in St. Louis

Citation: Second scientific balloon launches from Antarctica (2019, January 2) retrieved 18 April 2024 from <https://phys.org/news/2019-01-scientific-balloon-antarctica.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.