

# Scientists nail down important plant compound pathway

January 25 2019, by Brian Wallheimer

---



Natalia Dudareva and colleagues identified a complete alternative pathway for the production of phenylalanine in plants. The compound is a precursor to others that are thought to have uses as flavors, fragrances, biofuels, pharmaceuticals and insecticides. Credit: Purdue University

Purdue University plant molecular biochemist Natalia Dudareva and

colleagues have described a complete second pathway used by plants to produce phenylalanine, a compound important for all living organisms.

Phenylalanine is a building block of proteins and the precursor to thousands of compounds. Plants use these compounds as aromatics to attract pollinators, for defense, reproduction, growth and development. Humans must get [phenylalanine](#) from plants, and they also use [natural products](#) derived from this compound in flavors, fragrances, biofuels, insecticides and pharmaceuticals.

It had long been accepted that plants employ one biological pathway to produce phenylalanine, but this did not explain all observed production of the compound. Dudareva, a Purdue distinguished professor in the Department of Biochemistry and researcher in the Purdue Center for Plant Biology, recently discovered the final step in the alternative pathway. Now she, along with graduate student Yichun Qian and research scientist Joseph Lynch from her lab, and collaborators John Morgan and his student Longyun Guo from Purdue's Davidson School of Chemical Engineering, have identified remaining steps, completing the picture.

The key finding, reported in the journal *Nature Communications*, is that the entire process happens in the cytoplasm rather than cell organelles called plastids.

"It splits completely at a different point than predicted, and the whole pathway is localized in cytoplasm. This was unexpected," Dudareva said. "The gene responsible for the committed step has been known for 20 years and was never, until now, considered to be involved in production of phenylalanine."

Phytochemicals derived from phenylalanine are thought to have uses for a broad range of cancer treatments, as well a pain management,

depression and Parkinson's disease, among others. Obtaining these compounds from [plants](#) has been difficult because of several tightly regulated steps that occur when the phenylalanine precursor is made in the plastids of cells. By uncovering this second pathway in cytoplasm, which is less stringently controlled, biotechnological methods for enhancing yield of phenylalanine can be developed.

"A next step will be to use this newly discovered pathway with relaxed regulation to produce more phenylalanine and phenylalanine-derived [compounds](#), which we may explore in future," Dudareva said.

**More information:** Yichun Qian et al. Completion of the cytosolic post-chorismate phenylalanine biosynthetic pathway in plants, *Nature Communications* (2018). [DOI: 10.1038/s41467-018-07969-2](https://doi.org/10.1038/s41467-018-07969-2)

Provided by Purdue University

Citation: Scientists nail down important plant compound pathway (2019, January 25) retrieved 24 May 2024 from <https://phys.org/news/2019-01-scientists-important-compound-pathway.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--