



Scientists identify toxic antipredator defense mechanism in locusts

January 28 2019


A PAN distribution in the body of gregarious nymphs



B PAN is produced in gregarious locusts



C Behavioral assays of gregarious nymphs to different dosages of PAN



(A) Whole-body PAN production by a gregarious locust. (B) PAN is present as the most abundant component in gregarious locusts, but is absent in solitary locusts. (C) Total time spent (s) of a fourth-instar gregarious locust in the PAN zone versus in the paraffin oil control zone over 10 min in the dual-choice bioassay Credit: IOZ




A team of scientists led by Prof. Kang Le at the Institute of Zoology, Chinese Academy of Sciences, has reported an unprecedented animal defense mechanism by which an olfactory aposematic (warning) signal can be converted to a hypertoxic chemical to facilitate an antipredator defense in locusts.

The findings were published in an article titled "Phenylacetonitrile in locusts facilitates an antipredator [defense](#) by acting as an olfactory aposematic signal and cyanide precursor" in *Science Advances* on Jan. 23, 2019.

Animal aggregation is a very common phenomenon with important biological significance in the natural world. High-density locusts can form large swarms that are at high risk of exposure to predators. Therefore, an effective defense strategy by gregarious locusts is crucial for their survival and migration.

A previous study showed that gregarious locusts may use warning coloration associated with toxic extracts from food plants for antipredator defense against predation. However, the functional coordination and biosynthesis of aposematic signals and toxins in locusts are poorly understood.

Using [chemical analysis](#), researchers revealed that phenylacetonitrile (PAN) is a dominant density-dependent volatile compound that can be detected in all tissues and body fluids of gregarious locusts (Fig. 1A), but is absent in solitary locusts (Fig. 1B). Behavioral assays excluded PAN role in [locust](#) aggregations (Fig. 1C).




(A) In a dual-choice bioassay conducted with paired locusts, the percentage (%) of the great tits selected and fed on hexane-treated solitary locusts (control) or PAN-treated solitary locusts. (B) First choice, injury rate, and consumption rate for the paired dsGFP- and dsCYP305M2 (dsRNAi)-injected gregarious locusts. (C) Selections of great tits between a hexane-treated dsCYP305M2-injected locust (control) and a PAN-treated dsCYP305M2-injected locust. Credit: IOZ

Using transcriptome analysis, RNA interference-mediated gene knockdown, and chemical analysis of deuterium-labeled compounds, the researchers identified for the first time that a novel cytochrome P450 (CYP) gene CYP305M2 catalyzes the oximation reaction in the biosynthesis pathway from phenylalanine to PAN in animal species.

They showed that this gene encodes a rate limiting enzyme whose [expression levels](#) directly respond to the changes in locust population density.

To test whether PAN production by locusts influences predation by the [great tit](#), the researchers performed a series of dual-choice and predation tests involving locusts with or without PAN load. They found that great tits refused to attack and feed on PAN-treated solitary locusts (Fig. 2A).



A cytochrome (CYP) gene (CYP305M2) that catalyzes the formation of (Z)-PAOx, a precursor in PAN and HCN biosynthesis from phenylalanine in gregarious locusts, was identified. CYP305M2 is present at barely detectable levels in solitary locusts. Aposematism with PAN can warn and deter consumers from poisoning by HCN generated from PAN. The yellow marker in gregarious locusts denotes the expression of CYP305M2. Against a green background, solitary locusts utilize to predators and are otherwise readily discovered and predated. Credit: IOZ

Further, they showed that dsCYP305M2-injected locusts are frequently attacked and are preferentially consumed by great tits over dsGFP-injected (Figure 2B) or PAN-treated dsCYP305M2-injected locusts (Fig. 2C). These findings suggest that PAN produced by gregarious locusts reduces bird predation and serves as an olfactory aposematic signal in locust defense.

Although biosynthesis and administration of hydrogen cyanide (HCN) is prevalent among plants, it is relatively rare in animals. Here, the

researchers revealed for the first time that a de novo biosynthetic pathway of HCN is present in locusts and HCN from the PAN precursor is significantly promoted when locusts are threatened by bird predators (Fig. 3). Thus, the plasticity of PAN biosynthesis in response to population density is crucial for the optimization of the antipredator defense strategies of locusts under diverse environmental conditions.

The results changed the previous viewpoint on the role of PAN in locust aggregations and provided solid evidence to support its role in locust defense. The outcome of this study is relevant not only to the deep understanding of prey-predator interactions, but also to the development of novel strategies to control outbreaks of agricultural pests.

More information: Jianing Wei et al, Phenylacetonitrile in locusts facilitates an antipredator defense by acting as an olfactory aposematic signal and cyanide precursor, *Science Advances* (2019). [DOI: 10.1126/sciadv.aav5495](https://doi.org/10.1126/sciadv.aav5495)

Provided by Chinese Academy of Sciences

Citation: Scientists identify toxic antipredator defense mechanism in locusts (2019, January 28) retrieved 25 April 2024 from <https://phys.org/news/2019-01-scientists-toxic-antipredator-defense-mechanism.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.