

Study says seas may be rising faster than thought

January 30 2019, by Barri Bronston



Fifth-year PhD student Molly Keogh shot this photo near Bohemia, looking northeast over the marshes of Breton Sound in southeastern Louisiana. Credit: Tulane University

A new Tulane University study questions the reliability of how sea-level

rise in low-lying coastal areas such as southern Louisiana is measured and suggests that the current method underestimates the severity of the problem. This research is the focus of a news article published this week in the journal [Science](#).

Relative [sea-level rise](#), which is a combination of rising water level and subsiding land, is traditionally measured using tide gauges. But researchers Molly Keogh and Torbjörn Törnqvist argue that in coastal Louisiana, tide gauges tell only a part of the story.

Tide gauges in such areas are anchored an average of 20 meters into the earth rather than at the ground surface. "As a result, tide gauges do not record [subsidence](#) occurring in the shallow subsurface and thus underestimate rates of relative sea-level rise," said Keogh, a fifth year Ph.D. student and lead author of the study.

"This study shows that we need to completely rethink how we measure sea-level rise in rapidly subsiding coastal lowlands" said Törnqvist, Vokes Geology Professor in the Tulane School of Science and Engineering.

The study, published in the open-access journal [Ocean Science](#), says that while tide gauges can accurately measure subsidence that occurs below their foundations, they miss out on the shallow subsidence component. With at least 60 percent of subsidence occurring in the top 5 meters of the sediment column, tide gauges are not capturing the primary contributor to relative sea-level rise.

An alternative approach is to measure shallow subsidence using surface-elevation tables, inexpensive mechanical instruments that record surface elevation change in wetlands. Coastal Louisiana already has a network of more than 300 of these instruments in place. The data can then be combined with measurements of deep subsidence from GPS data and

satellite measurements of sea-level rise, Keogh said.

Rates of relative sea-level rise obtained from this approach are substantially higher than rates as inferred from tide-gauge data. "We therefore conclude that low-elevation coastal zones may be at [higher risk](#) of flooding, and within a shorter time horizon, than previously assumed," Keogh said.

She said the research has implications for [coastal communities](#) across the globe.

"Around the world, communities in low-lying [coastal areas](#) may be more vulnerable to flooding than we realized. This has implications for coastal management, city planners and emergency planners. They are planning based on a certain timeline, and if sea level is rising faster than what they are planning on, that's going to be a problem."

More information: Molly E. Keogh et al, Measuring rates of present-day relative sea-level rise in low-elevation coastal zones: a critical evaluation, *Ocean Science* (2019). [DOI: 10.5194/os-15-61-2019](https://doi.org/10.5194/os-15-61-2019)

Provided by Tulane University

Citation: Study says seas may be rising faster than thought (2019, January 30) retrieved 27 April 2024 from <https://phys.org/news/2019-01-seas-faster-thought.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.