

Long term ag change impacts stream water quality

January 2 2019



Graduate students Heather Luken and Tanner Williamson collect water samples from an automated sampler on Four Mile Creek. The autosampler collects a water sample from the stream every seven hours, and stores 24 samples in the carousel of bottles shown here. Credit: Carrie Ann Sharitt

In the early 1990s, Acton Lake in southwestern Ohio had a muddy

problem. Large amounts of sediment from nearby farms were entering the lake's watershed. These sediments traveled through streams draining the landscape and were filling up the lake.

So, the USDA gave local farmers incentives to change some of their farming practices. One of these practices was conservation tillage, in which the soil is plowed less often. That can reduce sediment runoff.

A new study examines how the switch to conservation tillage has impacted Acton Lake over the past decades. From 1994 to 2014, the researchers measured concentrations of suspended sediment, nitrogen, and phosphorus in streams draining into Acton Lake.

"We find that short-term trends in water quality may not reflect long-term changes," says study co-author Michael Vanni.

Tracking changes in water quality over the long term is vital, says Vanni, a biologist at Miami University, Ohio. "We don't have a lot of long-term information on how water quality in a stream or lake responds to agricultural change," he says.

That might be surprising since many ecologists study agricultural watersheds. But according to Vanni, studies on a given ecosystem are usually short term. "Long-term studies, like ours, can reveal important shifts in water quality," says Vanni. "Many of the changes we observed can only be seen after studying the streams for 20 plus years."

Vanni and his colleagues found that water quality responses were different during the first decade of the study (1994-2003) compared to the next (2004-2014). They also discovered that concentrations of suspended sediment, nitrogen, and phosphorus each reacted differently.

Levels of suspended [sediment](#) declined throughout the entire study

period. However, the decline was much sharper in the first ten years.

Phosphorus and nitrogen levels had contrasting outcomes. "The concentration of dissolved phosphorus in the streams declined sharply during the first ten years," says Vanni. "But then, phosphorus levels increased over the next ten years."



The inside of the automated sampler on Four Mile Creek, with the carousel of bottles used to collect water samples. The PVC pipe to the left houses a pressure transducer connected to a data logger and records the stage (height) of the stream water every ten minutes for a continuous record of stream flow. Credit: Carrie Ann Sharitt

In contrast, nitrogen levels didn't change much in the first ten years. After that, they fell sharply.

The study focused on the watershed of the Upper Four Mile Creek, which drains into Acton Lake. Most of the surrounding area is made up of corn and soybean farms. The researchers have monitored [farming practices](#) in the area since 1989 and water quality since 1994.

The long-term changes seen in this study indicate that there might be tradeoffs in managing different aspects of water quality. "The main reason to encourage [conservation tillage](#) was to reduce soil erosion and sedimentation in Acton Lake," says Vanni. "That has clearly been successful. Sediment inputs to the lake have declined."

Nitrogen levels are also declining. "That's great for local freshwater ecosystems," says Vanni. "It's also beneficial to the Gulf of Mexico, where some of our runoff eventually travels."

On the other hand, rising phosphorus levels are a cause for concern. "They could promote [algal blooms](#) downstream," says Vanni. "We might need to consider the tradeoffs involved in managing for sediments, nitrogen, or phosphorus."

It's not completely clear how the study findings would apply to other areas. However, the changes in [water](#) quality observed in this study are similar to those seen in some of the rivers that drain into Lake Erie.

High phosphorus levels are a problem in those watersheds as well. In fact, "high levels of phosphorus are implicated in causing increased blooms of harmful algae in Lake Erie," says Vanni.

Vanni and colleagues hope to continue measuring changes in suspended sediments, [nitrogen](#), and [phosphorus](#) in the Acton Lake watershed.

"We are also looking at the response of the Acton Lake ecosystem to these changes," says Vanni. "It will be really interesting to observe these changes over the next ten years, both from a scientific perspective as well as for [water quality](#) management."

More information: William H. Renwick et al, Stream Nitrogen, Phosphorus, and Sediment Concentrations Show Contrasting Long-term Trends Associated with Agricultural Change, *Journal of Environment Quality* (2018). [DOI: 10.2134/jeq2018.04.0162](https://doi.org/10.2134/jeq2018.04.0162)

Provided by American Society of Agronomy

Citation: Long term ag change impacts stream water quality (2019, January 2) retrieved 13 August 2024 from <https://phys.org/news/2019-01-term-ag-impacts-stream-quality.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.