

New theory sends temperatures to new lows

January 25 2019



Credit: University of Nottingham

Researchers have developed a new theory for recording the lowest temperatures ever measured, with the largest accuracy allowed by the laws of Nature. This line of research holds promise to revolutionise lowtemperature physics and could find a plethora of applications in emerging quantum technologies.

A recent collaboration involving a team from the University of Nottingham and The Institute of Photonic Sciences (Barcelona, Spain), shows that it is possible, in principle, to measure temperatures below a billionth of a Kelvin (!) in a cold atomic gas without disturbing it significantly, beating current precision standards. The work has been published in the latest edition of the journal *Physical Review Letters*.

In their study, the researchers modelled a Bose–Einstein condensate—a unique state of matter achieved by cooling an atomic gas down to extremely low temperatures—using realistic experimental parameters.



The thermometric technique would work by embedding an impurity atom into the atomic condensate, so that it acquires information about the <u>temperature</u> of the sample through interaction. In particular, its position and velocity become temperature-dependent so that, by monitoring them, the temperature can be inferred with high accuracy without disturbing the condensate.

Cooling atomic gases

Ultra-cold atomic gases are a very versatile experimental platform for a number of applications such as simulation of strongly correlated systems, quantum information processing, or the production of high-quality (cold) electron beams for electron microscopy or electron diffraction. For most of these applications it is essential to cool down the atomic gas to the lowest temperatures possible. Determining the temperature of these systems precisely is also critical for applications.

Mohammad Mehboudi, the lead author of the paper said: "The most common thermometric techniques currently available for cold atoms are destructive; that is, the sample is destroyed as a result of the measurement. On the other hand, non-destructive techniques usually lack the necessary accuracy at very low temperature. Our research provides a solution that overcomes both of these problems".

Outstanding experimental achievements allow nowadays high precision thermometry at very low temperatures. However, depending on the specific experimental platform, the underlying physical mechanism, accuracy, and effective temperature range of different thermometric schemes varies appreciably. Dr. Luis Correa also worked in the study and points out: "The newly-developed theoretical framework of quantum thermometry seeks to determine the fundamental limits on the precision of temperature measurements close to absolute zero; and it applies universally to any system. Importantly, this can provide clues as to how



to improve current low-temperature thermometric standards."

More information: Mohammad Mehboudi et al. Using Polarons for sub-nK Quantum Nondemolition Thermometry in a Bose-Einstein Condensate, *Physical Review Letters* (2019). DOI: 10.1103/PhysRevLett.122.030403

Provided by University of Nottingham

Citation: New theory sends temperatures to new lows (2019, January 25) retrieved 15 August 2024 from <u>https://phys.org/news/2019-01-theory-temperatures-lows.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.