

How trees and turnips grow fatter

January 9 2019



Root meristem (procambium) and 10-days-old root (active cambium). Credit: Pawel Roszak

Two international research teams have identified key regulatory networks controlling how plants grow 'outwards', which could help us to grow trees to be more efficient carbon sinks and increase vegetable crop yields.

As we crane our necks to stare up in awe at the height of the world's



tallest trees, it's easy to forget that what allows these living skyscrapers to stand for hundreds and even thousands of years is not upward growth, but outward growth, which makes stems and trunks fatter. Radial growth provides physical support to <u>plants</u>, yields everyday items like wood and cork, and plays a major role in converting atmospheric carbon into plant biomass.

Radial growth also produces specialised vascular tissues that transport water and nutrients around plants and is visible as concentric patterns known as annual growth rings in tree trunk cross-sections. In many cases, plants and trees continue this outwards growth for their entire lifetime. Radial growth is also responsible for producing our root and tuber vegetables such as turnips, carrots, sugar beet and potatoes. Yet the mechanisms behind how plants thicken their stems and roots by growing radially are virtually unknown.

Plant scientists from the University of Helsinki and University of Cambridge have independently uncovered two regulatory networks directing this all-important radial growth in plants. Their discoveries, published today in two separate papers in *Nature*, provides the most comprehensive understanding yet of how plants grow radially. It also means that botany textbooks around the world will need to update their descriptions of how the two types of vascular tissues in plants—xylem (better known as wood) that transports water and phloem that transports nutrients—form, differentiate and grow.





PEAR 1 protein (yellow) moving out from the phloem cell file (red). Credit: Pawel Roszak

Understanding how plants control and direct radial growth

Using thale cress (*Arabidopsis thaliana*) as a model plant to study how cell patterning develops during radial growth, Dr. Ari Pekka Mähönen's team from the Institute of Biotechnology (a HiLIFE unit) and Faculty of Biological and Environmental Sciences at the University of Helsinki has ended a near 150-year-old debate on the location and essence of the <u>stem</u> cells responsible for radial growth. His team showed that the stem cells for cambium (the tissue layer that gives rise to secondary stage vascular tissue) are located next to the young xylem. Furthermore, he showed that young xylem acts as an organiser for the stem cell activity.

Using the same model plant, Professor Ykä Helariutta's team at the



Sainsbury Laboratory Cambridge University (SLCU) focused on the early (primary) stage of vascular development. They showed that in contrast to the late stage, during this early stage young phloem cells (protophloem) are initiating and organising the primary (procambial) stage of radial growth. They also described an underlying gene regulatory network and an integrating role for a newly identified group of mobile transcription factors.

Together, their findings reveal some of the regulatory mechanisms that enable plants to continue to radially grow in a highly organised fashion, resulting in the concentric patterns seen in cross-sections of stems and roots. The key is the positioning of cells and a complex network of feedback signalling.

Dr. Mähönen's team combined individual cell lineage tracing and molecular genetics to show early-stage xylem cells, which had not yet differentiated, take over as the organiser and direct adjacent vascular cells to divide and function as stem cells: "We showed that this secondary development is a tightly controlled process and revealed a dynamic nature of the organiser. Differentiation of the organiser into a xylem vessel leads to formation of a new organiser in the adjacent cambial stem cell, thus ensuring the maintenance of the vascular cambium. We also identified a molecular mechanism that defines the stem cell organiser."

What is the nature of the intercellular signals that regulate vascular meristem activity?

Radial growth by vascular cambium represents the late, secondary stage of plant development particularly evident in tree species. The radial growth is primed in early seedlings during the so-called procambial stage, which is observed in all plant species.



"Very early-stage phloem tissues (protophloem) are helping to guide the behaviour of cells and establish patterns of developmental potential that impact on future radial growth," says Professor Helariutta. "This is established through a group of mobile transcription factors that move from the protophloem sieve element (PSE) to the neighbouring <u>cells</u>, to promote cell division, and to develop their own identity. The activity of these mobile transcription factors is regulated by a set of signals of various chemical nature, such as plant hormones, other transcription factors and mobile microRNA species."

Dr. Mähönen's and Professor Helariutta's teams are continuing to decipher how other cambial factors regulate the signalling network defining the stem cell organiser and how early stage radial growth influences secondary radial growth, respectively. This understanding could assist with future breeding of agricultural crops and trees to obtain higher economic yields and to maximise atmospheric carbon sequestration by increasing forest biomass.

More information: Shunsuke Miyashima et al, Mobile PEAR transcription factors integrate positional cues to prime cambial growth, *Nature* (2019). DOI: 10.1038/s41586-018-0839-y

Provided by University of Cambridge

Citation: How trees and turnips grow fatter (2019, January 9) retrieved 3 May 2024 from <u>https://phys.org/news/2019-01-trees-turnips-fatter.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.