


3,000-year-old eastern North American quinoa discovered in Ontario

January 15 2019



The monochrome shot is two seeds stuck together viewed from the side. This

shows the thin seed coats closely fitting the shape of the starchy endosperm inside the seed. Credit: Gary Crawford

A mass of charred seeds found while clearing a home construction site in Brantford, Ontario, has been identified as ancient, domesticated goosefoot (*C. berlandieri* spp. *jonesianum*), a form of quinoa native to Eastern North America. The seeds date back to 900 B.C., and have never previously been found north of Kentucky this early in history, says Professor Gary Crawford of the Department of Anthropology at the University of Toronto Mississauga (UTM), who was brought in by Archaeological Services Inc. (ASI), the archaeological consulting firm that excavated the site.

Archaeological discoveries don't normally shock Crawford but this one comes close. "Finding domesticated seeds that are so old in Ontario is special," Crawford says. "The next time we find a crop in the province is about 500 A.D., and it's corn. All previous research on this species of quinoa, which is now extinct, has taken place in the central United States: Arkansas, Illinois and Kentucky."

The charred seeds, about 140,000 in total, were discovered in Brantford in 2010 during a required archeological assessment conducted by Archaeological Services Inc. prior to site development. The Tutela Heights site, which has since become a housing development, yielded some [stone tools](#), post holes, debris and the chenopod seeds. Jessica Lytle, a co-author of the resulting [research paper](#), was one of the assessors who did the initial [seed](#) analysis and brought them to Crawford for further analysis, having studied with him at UTM. Their findings are published in the October 2018 issue of *American Antiquity*. The analysis took time, especially given the number of seeds and the need to document whether the whole collection was from the same crop.

"This discovery raises more questions than it answers. We had to consider whether the seeds were only traded here or grown locally," says Ron Williamson, Ph.D., of ASI, another co-author. "We also had to consider whether this was the beginning of agriculture in the province. It appears not, because we don't see any evidence of local cultivation. If it were grown in the region, we would have expected to see seeds of the crop in other pits around the site, but they were confined to this specific pit. We also don't see any sign of agricultural weeds or stone tools that may have been used for cultivation.

Indigenous peoples at the time exchanged certain kinds of minerals and finished stone objects over long distances, but this is the first evidence of a crop circulating in this exchange system. What meaning this plant had for local indigenous people nearly 3000 years ago still is not clear.



The color seed shot shows the crop (left) and the wild/weedy relative (right).

Credit: Gary Crawford

Professor Crawford notes "We always wondered if they were also exchanging perishable materials. We're taking the conservative view that these seeds were traded; it would make sense that it wasn't only stone and minerals being moved around. In Kentucky, Illinois and Arkansas, this was a very important foodstuff; its nutrient value was probably similar to that of modern quinoa, which comes from South America."

The researchers also explored how and why the seeds were charred. They speculate that it may have happened accidentally when the local inhabitants were attempting to parch them.

"You can lightly parch seeds so they don't sprout and store them," Crawford says. "It could have been a mistake to have burned them. There was a slight oxidization of the surrounding sediment, so the soil was heated; we think they were burned in place in the pit."

For Crawford, the next step in answering some of the questions will be to review seeds in his lab that were collected at other sites in Ontario to see if there are other charred seeds that may be variations of this subspecies and to examine other Ontario seed collections. Today, there is a weedy version that grows locally and he is curious whether this is a holdover from Indigenous agriculture.

"All of these bits of data demonstrate that the Indigenous Canadians were knowledgeable, sophisticated and well-connected across Eastern North America," Crawford says.

More information: Gary W. Crawford et al, AN EARLY WOODLAND DOMESTICATED CHENOPOD (CHENOPODIUM

BERLANDIERI SUBSP. JONESIANUM) CACHE FROM THE TUTELA HEIGHTS SITE, ONTARIO, CANADA, *American Antiquity* (2018). [DOI: 10.1017/aaq.2018.75](https://doi.org/10.1017/aaq.2018.75)

Provided by University of Toronto

Citation: 3,000-year-old eastern North American quinoa discovered in Ontario (2019, January 15) retrieved 20 April 2024 from <https://phys.org/news/2019-01-year-old-eastern-north-american-quinoa.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.