

# 400,000 African pangolins are hunted for meat every year—why it's time to act

February 14 2019, by Daniel J Ingram

---



Credit: AI-generated image ([disclaimer](#))

Pangolins, a group of unique African and Asian scaly mammals, are considered to be one of the most heavily trafficked wild mammals in the world. They are hunted and traded for their meat, scales, and other body parts, and used as traditional medicines in parts of Africa and Asia.

Of the eight pangolin species, [four](#) are found in Africa. These are the white-bellied, black-bellied, giant, and Temminck's ground pangolin. Three of these species live in Central African forests. The tree-dwelling white-bellied and black-bellied pangolins, weighing approximately 1.5 to 3kg (comparable to a small rabbit), and the ground-dwelling giant pangolin can weigh up to 33kg (the weight of a small Labrador dog).

But little is known about population sizes, mortality rates, and reproductive potential of African pangolins. Mounting evidence [suggests](#) that as the availability of Asian pangolins declines, and [international trade](#) flows increase, traders increasingly supply the more abundant and less expensive African pangolins to meet demand.

Seizures of pangolins and their scales and skins from Africa, destined for Asia, are [increasing](#) with [over](#) 53 tons seized in 2013 alone. These estimates likely represent a fraction of all pangolins traded, and an even smaller portion of the number of pangolins hunted.

To better understand how many pangolins are hunted in Central Africa each year, I and a team of researchers [collated](#) information on the quantities of animals that hunting villages extract from the forest, from studies conducted over the last 20 years. By doing this we can provide crucial information on regional trends which can be used to inform conservation actions and policy.

## **What we found**

We looked into the number of animals hunted in villages and offered for sale by collating data from research and reports that covered over 100 areas in sub-Saharan Africa between 1975 and 2014.

We extracted information on whether the animal was eaten or sold, how they were hunted, the sex, age category, and price. Other species

typically hunted for meat include blue duikers, brush-tailed porcupines and greater cane rats.

For Central African forests in Cameroon, Central African Republic, Equatorial Guinea, Gabon, Democratic Republic of Congo and Republic of Congo, we estimated that at least 400,000 pangolins are hunted annually for meat.



Seized pangolin scales from Cameroon. Credit: Linh Nguyen Ngoc Bao/MENTOR-POP

But we don't yet know whether pangolins can withstand these levels of hunting. This is mainly because we don't yet have reliable pangolin

population estimates for any of the species that inhabit Central African forests. Ideally, we would also need population and hunting data in the same location to be able to understand the levels of hunting that lead to population declines.

## **Pressures**

The pressures on African pangolins are likely increasing for several reasons.

Firstly, [increasing deforestation](#) across West and Central African countries has reduced their habitat, particularly for the semi-arboreal white-bellied pangolin and the arboreal black-bellied pangolin, which rely on forest habitats.

As the [human populations grow](#) in West, Eastern and Central Africa, this may exacerbate trends in deforestation and wildlife consumption.

Secondly, increases in the [accessibility](#) of remote areas to people and extractive industries may lead to more pangolin hunting. For example, a recent [study](#) showed that Asian industry workers in Gabon requested pangolins from hunters more than any other species.

Finally, the international trafficking of pangolins over the past decade has boomed. They are one of the most trafficked wild mammals in the world. For example, eight tonnes of pangolin scales trafficked from Nigeria, one of the largest ever hauls of scales, was [intercepted](#) a couple of weeks ago in Hong Kong.

## **Time to act**

While the media has greatly [increased](#) its coverage of the plight of



pangolins in recent years, financial and political support for conservation is still greatly needed. This includes support for pangolin population monitoring, identification of [pangolin](#) strongholds and areas in need of conservation, and the identification, design and testing of conservation interventions, where needed.

Without these steps we may see the African pangolins follow in the footsteps of their Asian counterparts.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: 400,000 African pangolins are hunted for meat every year—why it's time to act (2019, February 14) retrieved 30 April 2024 from <https://phys.org/news/2019-02-african-pangolins-meat-yearwhy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.