

Antibiotic resistances spread faster than so far thought

February 18 2019



Piaractus mesopotamicus, a South American species known as pacu, is often raised in aquaculture. Credit: Helmholtz Zentrum München



By studying fish raised in aquaculture, researchers from the Helmholtz Zentrum München, the University of Copenhagen and the University of Campinas in Brazil have shed new light on the mechanisms by which antibiotic resistance genes are transferred between bacteria. According to the study published in the journal 'Microbiome', those mechanisms are more varied than previously thought.

"In the past 70 years, the use of <u>antibiotics</u> in human and <u>veterinary</u> <u>medicine</u> has steadily increased, leading to a dramatic rise in resistant microorganisms," says Prof. Dr. Michael Schloter, head of the Research Unit for Comparative Microbiome Analyses (COMI) at Helmholtz Zentrum München. It is especially alarming that many microorganisms are resistant not just to one antibiotic, but to a whole range of different substances, states the corresponding author of the recent study. This poses particular problems in the treatment of infectious diseases. "We therefore set out to discover the mechanisms responsible for resistance development," he says.

To this end, he and his team, together with Danish scientists led by Gisle Vestergaard (University of Copenhagen and Helmholtz Zentrum München), investigated <u>fish</u> raised in aquaculture. Specifically, they studied Piaractus mesopotamicus, a South American species known as pacu that is often raised in aquaculture. The fish received the antibiotic florfenicol in their food for 34 days. During this time and after the application period, the researchers took samples from the digestive tract of the fish and looked for relevant genetic changes in the gut <u>bacteria</u>.

Resistance genes hop around the genome

"As expected, administration of the antibiotic induced an increase in the <u>genes</u> responsible for resistance to that antibiotic," explains COMI doctoral student Johan Sebastian Sáenz Medina, lead author of the paper. "One example are genes for pump proteins, which simply remove the



active substance from the bacteria again. However, we were particularly surprised by the different mechanisms that we could detect by which <u>antibiotic resistance genes</u> are spread amongst gut bacteria of the fish" Sáenz Medina explains. "This suggests that the bacteria also exchange resistance through viruses, known as phages, and transposons."

Further metagenomic studies confirmed that these mobile genetic elements induce a fast distribution of resistance genes among genomes of different organisms. So far it has been postulated that only plasmids (in essence, easily transferable mini-chromosomes) are mainly responsible for the exchange of resistance genes.

"The finding that <u>resistance</u> is also extensively transferred between bacteria without the involvement of plasmids is really quite surprising," says Michael Schloter. "Based on this observation, relevant dissemination models should be reviewed and modified. In addition, our data certainly lead us to question whether and to what extent we should continue to use antibiotics in the world's increasing number of aquacultures."

More information: Johan S. Sáenz et al. Oral administration of antibiotics increased the potential mobility of bacterial resistance genes in the gut of the fish Piaractus mesopotamicus, *Microbiome* (2019). DOI: 10.1186/s40168-019-0632-7

Provided by Helmholtz Association of German Research Centres

Citation: Antibiotic resistances spread faster than so far thought (2019, February 18) retrieved 8 May 2024 from <u>https://phys.org/news/2019-02-antibiotic-resistances-faster-thought.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.