

Research forms complex picture of mercury pollution in a period of global change

February 14 2019



A school of sardines in Italy. Credit: Wikimedia / Alessandro Duci

Climate change and the loss of wetlands may contribute to increased levels of mercury concentrations in coastal fish, according to a Dartmouth College study.

The finding implies that forces directly associated with [global change](#)

—including increased precipitation and land use modifications—will raise levels of the toxic metal that enter the marine food chain.

Estuaries, including coastal wetlands, provide much of the seafood that is harvested for human consumption and also serve as important feeding grounds for larger marine fish.

The study, published in late December in the journal *Environmental Pollution*, adds to the mounting body of research that indicates a complex relationship between the environment and [mercury](#) pollution.

"Estuaries provide habitat for the fish that feed our families," said Celia Chen, director of the Dartmouth Toxic Metals Superfund Research Program. "It's important to understand how mercury acts within our environment, particularly under increasing climate and land use pressures."

The Dartmouth study concludes that higher levels of mercury, and its toxic form [methylmercury](#), are associated with higher organic carbon in coastal waters. The study also finds that this results in higher levels of mercury occurring in fish that frequent these waters.

While the amount of mercury deposited from the atmosphere has decreased in parts of the U.S., elevated levels of the pollutant already accumulated in the soils from past deposition may be released into rivers and streams with the heavier rains associated with climate change.

At the same time, levels of organic carbon reaching estuaries in the region are also predicted to rise with rainfall. Since organic carbon binds with mercury, these increases may bring higher levels of mercury to estuarine waters.

The research is particularly relevant in the Northeast part of the country

given the high levels of existing mercury accumulated in the landscape and the increase in rainfall.

"Taken as a whole, these ecosystem changes have unpredictable effects on methylmercury bioaccumulation," said Vivien Taylor, a research scientist in the Department of Earth Science at Dartmouth and lead author of the study.

The Dartmouth study investigated methylmercury in intertidal zones located at three different latitudes and temperature regions in the Northeast region of the United States: Mount Desert Island, Maine; Long Island Sound, Connecticut and the upper Chesapeake Bay in Maryland.

The researchers studied mussels, periwinkle, shrimp, crabs and a variety of fish to confirm how carbon and methylmercury interact to enter the food chain in estuaries. In addition to chemical signatures, the team factored in parameters including fish size, tidal conditions and background levels of mercury contamination in regional sites.

The study helps untangle the competing processes that affect the levels of mercury that end up in fish.

"The interaction between climate change and mercury are complex and require much more study. There is, however, increasing evidence that some factors related to climate change might aggravate [mercury pollution](#)," said Chen.

The loss of coastal wetlands, further complicates the picture. While wetlands can be a source of methylmercury, organic carbon from wetlands may help reduce the amount of mercury that passes into fish and other aquatic species. The conversion of wetlands to other uses means that nature loses an important buffer that keeps mercury out of the food supply.

The study finds that organic carbon's ability to reduce the amount of methylmercury available to estuarine fish can be overwhelmed by the increased watershed inputs of dissolved organic carbon and methylmercury associated with [climate change](#). The result is increased methylmercury levels in estuarine species, like silversides and mummichog, which are food for larger [fish](#) like striped bass.

More information: V.F. Taylor et al, Organic carbon content drives methylmercury levels in the water column and in estuarine food webs across latitudes in the Northeast United States, *Environmental Pollution* (2018). [DOI: 10.1016/j.envpol.2018.12.064](https://doi.org/10.1016/j.envpol.2018.12.064)

Provided by Dartmouth College

Citation: Research forms complex picture of mercury pollution in a period of global change (2019, February 14) retrieved 28 April 2024 from <https://phys.org/news/2019-02-complex-picture-mercury-pollution-period.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--