

## A new method for precision drug delivery: painting

## February 27 2019, by Brandie Jefferson



Researchers from the McKelvey School of Engineering and Washington University School of Medicine in St. Louis are one step closer to delivering precise amounts of medication to exact location, repurposing an existing imaging 'painting' method. Cavitation images (left); PET images (middle); overlay of the two (right). (Courtesy: Hong Chen lab) Credit: Hong Chen lab



If traditional drug delivery were a type of painting, it might be akin to paintball. With good aim, a majority of the paint ends on the bullseye, but it also drips and splashes, carrying streams of paint across the target.

If the <u>drug</u> needs to enter the bloodstream and circulate throughout your body for treating disease wherever it may be, this paintball-like <u>delivery</u> <u>system</u> may work. But it won't work for targeted and precise drug delivery.

A more acute delivery approach would look more like "painting by numbers," a technique that would allow precise delivery of a certain amount of drugs to an exact location. Researchers at the James McKelvey School of Engineering and the School of Medicine at Washington University in St. Louis are developing the tools necessary for such a drug delivery system, which they call cavitation dose painting.

Their research was published online this week in Scientific Reports.

Using focused <u>ultrasound</u> with its contrast agent, microbubbles, to deliver drugs across the <u>blood-brain barrier</u> (FUS-BBBD), the research team, led by Hong Chen, assistant professor of biomedical engineering at McKelvey School of Engineering, and assistant professor of radiation oncology at the School of Medicine, was able to overcome some of the uncertainty of drug delivery.

This method takes advantage of the microbubbles expanding and contracting when they interact with the ultrasound, essentially pumping the intravenously-delivered drug to wherever the ultrasound is pointing.

To determine where and how much of the drugs were being delivered, the researchers used nanoparticles tagged with radio labels to represent drug particles, then used positron emission tomography (PET) imaging to track their whereabouts and concentrations. They could then create a



detailed image, showing where the nanoparticles were going and in what concentrations.

There's one hitch, though.

"The problem is, PET imaging is expensive and associated with radioactive exposure," Chen said.

So the team turned to passive cavitation imaging (PCI), an ultrasound imaging technique that has been under development by several groups for imaging the spatial distribution of microbubble cavitation, or the oscillation of microbubbles in the ultrasound field.

To determine whether PCI could also accurately determine the amount of drugs at a certain location, they correlated a PCI image with a PET image (which they knew can quantify the concentration of radioactive agents).

"We found there's pixel by pixel correlation between the ultrasound imaging and the PET imaging," said Yaoheng Yang, the lead author of this study and a second-year Ph.D. student in the Department of Biomedical Engineering. The PCI image, therefore, can be used to predict where a drug goes and how much drug is there. Hence, she called the new technique cavitation dose painting.

Going forward, Chen believes this method could drastically change the way some drugs are delivered. Using cavitation dose painting in tandem with focused ultrasound will allow doctors to deliver precise amounts of drugs to specific locations, for example, targeting different areas of a tumor with exactitude.

"I think this cavitation dose painting technique in combination with focused ultrasound-enabled brain drug delivery opens new horizons in



spatially targeted and modulated brain drug delivery," Chen said.

She has recently received a \$1.6 million grant from the National Institutes of Health (NIH)'s National Institute of Biomedical Imaging and Bioengineering to work on combining intranasal drug <u>delivery</u> and focused ultrasound (FUSIN) with this research.

The research team from Washington University School of Medicine included Xiaohui Zhang, postdoctoral research associate of radiology; Richard Laforest, associate professor of radiology; Yongjian Liu, associate professor of radiology; and Jeffrey F. Williamson, professor of radiation oncology. From McKelvey Engineering: Haoheng Yang; and Dezhuang Ye.

**More information:** Yaoheng Yang et al, Cavitation dose painting for focused ultrasound-induced blood-brain barrier disruption, *Scientific Reports* (2019). DOI: 10.1038/s41598-019-39090-9

## Provided by Washington University in St. Louis

Citation: A new method for precision drug delivery: painting (2019, February 27) retrieved 23 June 2024 from <u>https://phys.org/news/2019-02-method-precision-drug-delivery.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.