


Nanoparticles need some space to transfer energy

February 11 2019



Scientists are showing how energy moves (green) or doesn't (red) along nanoparticle chains. Credit: Journal of Materials Chemistry C. Royal Society of Chemistry

Quickly transferring energy from one place to another—without loss—could fundamentally change solar panels and computers. Materials made from long chains of tiny particles, or nanoparticles, show promise for such energy transfer. However, going even smaller than nano-dimensions hasn't worked quite as planned. Researchers suspected the particles needed to be close to each other to transfer energy efficiently. Now, a team has shown that the particles can be too close. Specifically, the energy transfer drops dramatically when the particles in the chain are less than a nanometer apart.

For those who want to build better [solar panels](#) or [computer chips](#), this study answers a basic question about the physics of a promising design. That design might use a [chemical structure](#) that contains chains of nanoparticles. The study explains why the [energy](#) transfer efficiency drops. That is, it shows how quantum mechanical effects alter the transfers. Also, it shows that complex calculations, using a real-time density functional tight binding approach, shed mechanistic insight to analyzing energy transfers based on interparticle distance.

Scientists wanted to understand the quantum-mechanical effects that can result in qualitatively different and sometimes completely opposite results. In particular, they wanted to understand the reason behind efficiencies and inefficiencies of energy transfer in long nanoparticle chains. Such chains hold promise in diverse fields, including energy harvesting. Conventional calculation approaches were not up to the task. Researchers used real-time density functional tight binding to characterize energy transfer as a function of interparticle distance. In [contrast](#) to classical electrodynamics methods, their quantum dynamical calculations show a drop in efficiency for spacings of subnanometer lengths within the nanoparticle chain. The drop in efficiency is due to quantum mechanical tunneling that dramatically changes the electronic couplings between nanoparticles. Thus, the team showed that considering nanoparticle spacing as well as larger quantum-mechanical effects is

vital to accurately calculate energy transfer mechanisms.

More information: Niranjan V. Ilawe et al. Effect of quantum tunneling on the efficiency of excitation energy transfer in plasmonic nanoparticle chain waveguides, *Journal of Materials Chemistry C* (2018). [DOI: 10.1039/C8TC01466C](https://doi.org/10.1039/C8TC01466C)

Provided by US Department of Energy

Citation: Nanoparticles need some space to transfer energy (2019, February 11) retrieved 25 April 2024 from <https://phys.org/news/2019-02-nanoparticles-space-energy.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.