

Organic particles modulate soot mixing in the atmosphere

February 13 2019



Credit: CC0 Public Domain

In the atmosphere, individual soot particles emitted from human activities often encounter and combine with organic material, forming soot-containing particles. A research team led by scientists from the U.S.

Department of Energy's (DOE) Pacific Northwest National Laboratory/EMSL and Michigan Technological University established a link between the viscosity of the organic material and its distribution within the soot-containing particle. This mixing configuration is important because it alters the particles' ability to absorb and scatter solar radiation.

The scientists discovered that organic material of low viscosity partially or completely encapsulated soot particles, while highly viscous organic material often attaches to the surface of co-existing soot particles or do not even stick together, remaining separate.

Soot particles strongly absorb solar radiation, and therefore, affect Earth's energy balance. However, their [optical properties](#) strongly depend on how the individual soot particles mix with other particles in the atmosphere (internal mixing).

This study identified the viscosity of organic matter in the atmosphere as a key factor that influences the distribution of organic matter within soot-containing particles. These findings will improve aerosol models that predict the mixing state of soot via condensation and coagulation processes, thereby improving radiative forcing calculations and climate predictions.

Organic particles in the atmosphere exhibit a wide range of viscosity, depending upon their [chemical composition](#) and the surrounding environmental conditions, such as temperature and relative humidity. As soot particles combine with organic particles in the atmosphere, the distribution of soot and organic material in the combined particle determines its overall optical properties and radiative forcing-warming or cooling effects.

Taking scanning electron microscopy images at different view angles,

researchers investigated soot-containing particles collected during DOE's Carbonaceous Aerosol and Radiative Effects Study (CARES) in Sacramento, California, in June 2010. Their results indicate that organic material of low to intermediate viscosity often engulfed soot, while highly viscous organic material attaches to the soot surface or remains separated from it. Soot particles engulfed by low viscosity organics become coated and can be approximately represented by a "core-shell" model, where the soot particles are at the center of an idealized spherical shell made of the organic material. In contrast, increasing viscosity results in limited (partial) engulfing, and hence the formation of non-core-shell configurations.

These findings establish a link between the viscosity of [organic material](#) and its distribution when mixed internally with [soot particles](#). Accounting for the effects of viscosity of organic matter on the mixing state of soot-containing [particles](#) in numerical models will improve estimates of radiative forcing from [soot](#).

More information: Noopur Sharma et al. Physical Properties of Aerosol Internally Mixed With Soot Particles in a Biogenically Dominated Environment in California, *Geophysical Research Letters* (2018). [DOI: 10.1029/2018GL079404](https://doi.org/10.1029/2018GL079404)

Provided by Environmental Molecular Sciences Laboratory

Citation: Organic particles modulate soot mixing in the atmosphere (2019, February 13) retrieved 26 April 2024 from <https://phys.org/news/2019-02-particles-modulate-soot-atmosphere.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.