


# **Sandia identifies unusual polycyclic aromatic hydrocarbons using tandem mass spectrometry**

February 7 2019, by Michael Padilla

---



Sandia National Laboratories scientists Scott Skeen, left, Nils Hansen, center, and Brian Adamson discuss tandem mass spectrometry, which was used for the detection of aliphatically linked polycyclic aromatic hydrocarbons found in soot producing flames. Credit: Michael Padilla

In most situations, breaking things apart isn't the best way to solve a problem. However, sometimes the opposite is true if you're trying to characterize complex chemical compounds. That's what Sandia National Laboratories scientists Nils Hansen and Scott Skeen did to definitively identify pollution-causing soot precursors in a flame.

The researchers discovered [polycyclic aromatic hydrocarbons](#) (PAHs) with aliphatic side chains, which have been hypothesized to serve as "seeds" for [soot](#) particles in [engine emissions](#).

"The role of these molecules as soot precursors has been hypothesized, and there is indirect experimental evidence of their presence on the surface of soot extracted from flames," said Skeen. "Until now, however, no one had definitive experimental proof of their existence as stable components in the flame."

The newly recognized compounds can be used to create more detailed models of combustion that, in turn, can help in the design of cleaner, more efficient engines that emit less soot and fewer harmful hydrocarbons into the atmosphere.

Working with former Sandia postdoctoral researcher Brian Adamson and Lawrence Berkeley National Laboratory's Musa Ahmed, Hansen and Skeen published the discovery in the *Journal of Physical Chemistry A*.

Funding for the research came from Sandia's Laboratory Directed Research and Development program, while Ahmed is supported by the Department of Energy's Office of Basic Energy Sciences.

The team used an analytical technique called tandem mass spectrometry—using an instrument provided by Lawrence Berkeley Lab and customized by Adamson—to detect gas-phase polycyclic aromatic hydrocarbons with aliphatic side chains in flames that produce soot, something never done before.

The device removes an electron to charge, or ionize, large molecules sampled from the flame, measures the molecules' masses then further identifies them by characterizing how the ionized molecules crack apart.

## **Latest discovery builds on previous Sandia research**


Recent work by Sandia scientist Hope Michelsen, technologist Paul Schrader and former postdoctoral researcher Olof Johansson broke ground by identifying chemical chain-reaction processes where hydrocarbons could form soot. That work heightened the challenge of detecting and characterizing the compounds that participate in these processes.

One area of debate is whether the chemical byproducts in soot are polycyclic aromatic hydrocarbons, made solely out of ring-shaped groups of atoms, or contain extra, non-cyclic structures called alkyl, or aliphatic, groups. These hydrocarbon chains can make the links among polycyclic aromatic hydrocarbons more stable at the high temperatures of combustion, greater than 3,600 degrees Fahrenheit.

"Without the tandem component of this new mass spectrometer, each molecule's mass is obtained but no information about its structure is revealed. You see something at mass 78, at mass 128, and so forth, but

you don't know which molecules are represented. You just use your chemical intuition," Hansen said. "Think of a mass spectrometer as an instrument that sorts a container full of mixed nuts based on the weight of each individual nut, but at the end you still don't know whether you sorted peanuts, hazelnuts or walnuts."

The customized tandem mass spectrometer that the team used makes it easier to characterize the structure of large molecules by breaking them apart through high-energy collisions in a collision induced dissociation cell.



A flame that was sampled by Sandia National Laboratories scientists against the backdrop of mass spectra and polycyclic aromatic hydrocarbon compounds found inside the flame. Credit: Sandia National Laboratories

"Normal mass spectrometry can tell you how many atoms of each

element are present in a molecule, but it won't tell you anything about how those atoms are joined together," Adamson said. "Tandem mass spectrometry with collision induced dissociation isolates molecules of a single mass and then breaks them apart. The way they break apart provides clues as to the structure of the parent molecule."

The team found direct evidence that aliphatically-bridged polycyclic aromatic hydrocarbons and PAHs with alkyl chains exist in the sampled gases of the soot-forming flame. Such species may be sufficiently stable at the high temperatures of combustion to serve as key components in incipient soot particle formation.

The team also used a special flame configuration to minimize disruptions to the flame chemistry caused by the sampling process. Skeen said the setup involved sampling and examining large molecules from an inverted candle-like flame.

"In a candle, the wax moves up the wick and then vaporizes before burning in the surrounding air. The flame appears yellow because soot particles get very hot as they move through the flame," Skeen said. "In this configuration, it is impossible to sample soot particles or molecules that lead to soot formation without disturbing the flame because a probe must be inserted through the flame sheet.

"To overcome this problem, we generated a flame in which the air is in the center of the flame with the fuel on the outside," he said. "This way, we can probe the gases of interest from the outside of this 'inverse' flame. This is perhaps the first time that such a [flame](#) has been attached to a tandem mass spectrometer."

## **Soot needs to be filtered for cleaner engines**

The search for soot precursors is motivated by the need for cleaner

engines that still run efficiently. Under certain driving conditions, diesel emissions exceed government regulations. This has led to the use of particulate filters that effectively capture [soot particles](#) from diesel exhaust, but they make the vehicles significantly more expensive and less efficient. Engines that produce less soot would need smaller particulate filters, reducing costs and increasing fuel economy.

Engine manufacturers typically use computer simulations to improve engine designs. They model the fuel injection, combustion and pollutant formation processes. Skeen said that better understanding of how soot compounds are produced—specifically definitive identification of polycyclic aromatic hydrocarbons with alkyl chains attached—should lead to models that more accurately describe the effects of engine design parameters on emissions and efficiency.

"If we can understand the chemistry, we can develop a model that will allow engine designers to optimize fuel injectors, air flows and the shape of internal engine surfaces, among other things, that will keep these compounds out of the atmosphere," Skeen said.

## **Future steps to develop new models for soot formation**

This discovery of alkyl-substituted and aliphatically-bridged polycyclic aromatic hydrocarbons in sooting flames is only the starting point for using tandem [mass spectrometry](#) to decipher the complex chemistry of polluting emissions, the team said.

Using this technique, thousands of different types of compounds could potentially be identified. Even for the most basic polycyclic aromatic hydrocarbons, there are about a hundred different ways the atoms can come together. Seeing all the different arrangements presents a

formidable challenge. Ahmed will continue his work with Sandia scientists and plans to introduce infrared spectroscopy for less ambiguous identification of alkyl-substituted and aliphatically-bridged PAHs in soot.

The Sandia scientists hope to collaborate with data scientists to develop more efficient, realistic models of engine soot formation, ultimately leading to designs for cleaner, more efficient engines.

**More information:** B. D. Adamson et al. Detection of Aliphatically Bridged Multi-Core Polycyclic Aromatic Hydrocarbons in Sooting Flames with Atmospheric-Sampling High-Resolution Tandem Mass Spectrometry, *The Journal of Physical Chemistry A* (2018). [DOI: 10.1021/acs.jpca.8b08947](https://doi.org/10.1021/acs.jpca.8b08947)

Provided by Sandia National Laboratories

Citation: Sandia identifies unusual polycyclic aromatic hydrocarbons using tandem mass spectrometry (2019, February 7) retrieved 20 June 2024 from <https://phys.org/news/2019-02-sandia-unusual-polycyclic-aromatic-hydrocarbons.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.