

Scientists find persistent organic pollutants in animal fur

February 14 2019



Hedgehog, Czech Republic. Credit: Tadeusz Lakota on Unsplash

Scientists of the Far Eastern Federal University (FEFU), working as part



of an international toxicologists' team, studied fur samples of wild terrestrial mammals in Primorye, Russia. All samples contained persistent organic pollutants (POPs) that are resistant to decomposition, tend to accumulate in body tissues and are potentially risky for human and animal health. Some of them are prohibited by the Stockholm Convention. The research outcome was published in *Environmental Science and Pollution Research*.

The highest POP content levels were found in the fur samples of musk deer, an <u>endangered species</u> listed in the Red Book. One of the reasons for such high toxicity could be the diet of the animals—musk deer feed on lichens, which accumulate <u>harmful substances</u> by absorbing them from the atmosphere.

In total, 15 animal hair samples were studied. They were isolated from six leopard cats, three musk deer, one Amur hedgehog, one wolf and four raccoons.

"In 73 percent of cases, we found traces of the insecticide hexachlorobenzene and DDT derivatives prohibited by the Stockholm Convention. In 100 percent of cases, <u>polycyclic aromatic hydrocarbons</u> (PAHs) were found, such as anthracene, fluoranthene, pyridine, and phenanthrene. PAHs are formed as a result of the thermal decomposition of cellulose, but most of these substances are of technogenic origin, i.e., they appear in the environment as a result of human activity. In most of the studied fur samples, we found at least 10 of the 16 types of POPs, the presence of which was the subject of our study. The study was innovative, as our international team found evidence of the accumulation of different POPs types in the hair of five species of wild terrestrial mammals at once," says Kirill Golokhvast, Ph.D., FEFU vice president for research.

The researchers point out that at the global level, the environmental



contamination by POPs compounds is increasing, which is of particular concern because they are harmful to humans, marine animals, and terrestrial ecosystems. Despite the fact that the global ban is imposed by the Stockholm Convention to the production and uses of such hazardous substances as hexachlorobenzene it continues to be found in living organisms throughout the world.

The researchers propose to further monitor the accumulation of POPs in the fur of a greater number of terrestrial mammals species because today the data on this topic are rather scarce.

Scientists consider the analysis of animal fur and human hair samples as effective, non-invasive and non-destructive methods of biological monitoring of ecosystem contamination. Animal fur samples are easy to collect, store and analyze. For the purpose of the study, the scientists took samples of the fur from the <u>animals</u> killed in Primorye as a result of traffic accidents or hunting.

More information: Evangelia I. Iatrou et al, Monitoring of environmental persistent organic pollutants in hair samples collected from wild terrestrial mammals of Primorsky Krai, Russia, *Environmental Science and Pollution Research* (2019). DOI: <u>10.1007/s11356-019-04171-9</u>

Provided by Far Eastern Federal University

Citation: Scientists find persistent organic pollutants in animal fur (2019, February 14) retrieved 3 May 2024 from <u>https://phys.org/news/2019-02-scientists-persistent-pollutants-animal-fur.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is



provided for information purposes only.