

Long term study shows sources of black carbon in the Arctic

February 14 2019, by Bob Yirka



The sampling hut in Tiksi. Credit: Oleg Dudarev


An international team of researchers has conducted the most thorough

study yet of the sources of black carbon in the Arctic. In their paper published in the journal *Science Advances*, the group describes their findings and explain why what they found relates to global warming.

While most of the press regarding [global warming](#) centers on the release of [carbon dioxide](#) into the atmosphere, very little is heard about other man-made activities that are also heating the planet. One such activity is the emission of [soot](#). It comes from factory chimneys, particularly those that burn coal, burning gasoline in car engines, burning fields as part of agricultural efforts, and from other activities that involve burning some type of material.

Because of inefficiencies, some bits of [carbon](#), in the form of soot, make their way into the atmosphere. They do not stay there nearly as long as carbon dioxide does—the problem occurs when they come back down. When it lands on ice in the Arctic, it turns the ice from white to black. Blackened ice absorbs heat, white ice reflects it. Thus, the carbon-covered ice tends to have more surface melting than normal, and there is much less reflection of heat back into the atmosphere. Both contribute to the rapidity of global warming happening in the Arctic compared to other parts of the world.

In recent years, several research efforts have explored the impact of black carbon in the Arctic and sought to ascertain its sources. The researchers with this new effort claim most prior studies have produced unreliable results due to what they describe as "a lack of observational restraints" and less-than-clear emission inventories. To address such problems, the team undertook a five-year study of black carbon in the Arctic, which involved collecting samples from multiple sites and subjecting each to [chemical analysis](#) to determine which kind of source it came from.



The Dr. Neil Trivett Global Atmosphere Watch (GAW) Observatory is the most northerly site in the global network, located at Alert, Nunavut, Canada (approximately 800km south of the geographic North Pole). Environment and Climate Change Canada operates the observatory, which hosts research campaigns and facilitates long-term measurements of a number of atmospheric constituents including greenhouse gases, aerosols including black carbon, stratospheric ozone, Persistent Organic Pollutants, mercury, etc. Credit: Janice Lange

The team reports that the majority of [black carbon](#) in the Arctic (approximately 70 percent) comes from burning [fossil fuels](#), not croplands (or wildfires) as some have suggested. But they also found that the ratios vary by season—during the winter, much more fossil fuel is burned to keep buildings warm. In contrast, most of the crop burning and

other types of fires occur in the warmer months.



Equipment being packed up for transport back to Stockholm. Credit: Patrik Winiger

More information: P. Winiger et al. Source apportionment of circum-Arctic atmospheric black carbon from isotopes and modeling, *Science Advances* (2019). [DOI: 10.1126/sciadv.aau8052](https://doi.org/10.1126/sciadv.aau8052)

© 2019 Science X Network

Citation: Long term study shows sources of black carbon in the Arctic (2019, February 14)
retrieved 3 May 2024 from

<https://phys.org/news/2019-02-term-sources-black-carbon-arctic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.