


Where is the universe hiding its missing mass?

February 15 2019



Credit: Chandra X-ray Center

Astronomers have spent decades looking for something that sounds like

it would be hard to miss: about a third of the "normal" matter in the Universe. New results from NASA's Chandra X-ray Observatory may have helped them locate this elusive expanse of missing matter.

From independent, well-established observations, scientists have confidently calculated how much normal matter—meaning hydrogen, helium and other elements—existed just after the Big Bang. In the time between the first few minutes and the first billion years or so, much of the normal matter made its way into cosmic dust, gas and objects such as stars and planets that telescopes can see in the present-day Universe.


The problem is that when astronomers add up the mass of all the normal matter in the present-day Universe about a third of it can't be found. (This missing matter is distinct from the still-mysterious [dark matter](#).)

One idea is that the missing mass gathered into gigantic strands or filaments of warm (temperature less than 100,000 Kelvin) and hot (temperature greater than 100,000 Kelvin) gas in intergalactic space. These filaments are known by [astronomers](#) as the "warm-hot intergalactic medium" or WHIM. They are invisible to optical light telescopes, but some of the warm gas in filaments has been detected in ultraviolet light.

Using a [new technique](#), researchers have found new and strong evidence for the hot component of the WHIM based on data from Chandra and other telescopes.

"If we find this missing mass, we can solve one of the biggest conundrums in astrophysics," said Orsolya Kovacs of the Center for Astrophysics | Harvard & Smithsonian (CfA) in Cambridge, Massachusetts. "Where did the universe stash so much of its matter that makes up stuff like stars and planets and us?"

Astronomers used Chandra to look for and study filaments of warm gas lying along the path to a quasar, a bright source of X-rays powered by a rapidly growing supermassive black hole. This quasar is located about 3.5 billion light years from Earth. If the WHIM's hot gas component is associated with these filaments, some of the X-rays from the quasar would be absorbed by that hot gas. Therefore, they looked for a signature of hot gas imprinted in the quasar's X-ray light detected by Chandra.



Light Path (Credit: NASA/CXC/K. Williamson, Springel et al.)

One of the challenges of this method is that the signal of absorption by the WHIM is weak compared to the total amount of X-rays coming from

the quasar. When searching the entire spectrum of X-rays at different wavelengths, it is difficult to distinguish such weak absorption features—actual signals of the WHIM—from random fluctuations.

Kovacs and her team overcame this problem by focusing their search only on certain parts of the X-ray light spectrum, reducing the likelihood of false positives. They did this by first identifying galaxies near the line of sight to the quasar that are located at the same distance from Earth as regions of warm gas detected from ultraviolet data. With this technique they identified 17 possible filaments between the quasar and us, and obtained their distances.

Because of the expansion of the universe, which stretches out light as it travels, any absorption of X-rays by matter in these filaments will be shifted to redder wavelengths. The amounts of the shifts depend on the known distances to the filament, so the team knew where to search in the spectrum for absorption from the WHIM.

"Our technique is similar in principle to how you might conduct an efficient search for animals in the vast plains of Africa," said Akos Bogdan, a co-author also from CfA. "We know that animals need to drink, so it makes sense to search around watering holes first."

While narrowing their search helped, the researchers also had to overcome the problem of the faintness of the X-ray absorption. So, they boosted the signal by adding spectra together from 17 filaments, turning a 5.5-day-long observation into the equivalent of almost 100 days' worth of data. With this technique they detected oxygen with characteristics suggesting it was in a gas with a temperature of about one million degrees Kelvin.

By extrapolating from these observations of oxygen to the full set of elements, and from the observed region to the local universe, the

researchers report they can account for the complete amount of missing matter. At least in this particular case, the missing matter had been hiding in the WHIM after all.

"We were thrilled that we were able to track down some of this missing matter" said co-author Randall Smith, also of CfA. "In the future we can apply this same method to other quasar data to confirm that this long-standing mystery has at last been cracked."

A paper describing these results was published in the *Astrophysical Journal* on February 13, 2019.

More information: Orsolya E. Kovacs et al. Detection of the Missing Baryons toward the Sightline of H1821+643. arXiv:1812.04625 [astro-ph.CO]. arxiv.org/abs/1812.04625

Provided by Chandra X-ray Center

Citation: Where is the universe hiding its missing mass? (2019, February 15) retrieved 27 January 2023 from <https://phys.org/news/2019-02-universe-mass.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.