


# Novel experiment validates widely speculated mechanism behind the formation of stars

February 5 2019, by John Greenwald

---



Water-filled version of MRI experiment showing transparent outer cylinder and blackened inner cylinder. Red lasers enter at bottom to measure the local speed of the water. Credit: Eric Edlund and Elle Starkman


How have stars and planets developed from the clouds of dust and gas that once filled the cosmos? A novel experiment at the U.S. Department of Energy's (DOE) Princeton Plasma Physics Laboratory (PPPL) has demonstrated the validity of a widespread theory known as "magnetorotational instability," or MRI, that seeks to explain the formation of heavenly bodies.

The theory holds that MRI allows [accretion disks](#), clouds of dust, gas, and plasma that swirl around growing stars and [planets](#) as well as black holes, to collapse into them. According to the theory, this collapse happens because turbulent swirling plasma, technically known as "Keplerian flows," gradually grows unstable within a disk. The instability causes angular [momentum](#)—the process that keeps orbiting planets from being drawn into the sun—to decrease in inner sections of the disk, which then fall into celestial bodies.

Unlike orbiting planets, the matter in dense and crowded accretion disks may experience forces such as friction that cause the disks to lose angular momentum and be drawn into the objects they swirl around. However, such forces cannot fully explain how quickly matter must fall into larger objects for planets and stars to form on a reasonable timescale.

## MRI experiment

At PPPL, physicists have simulated the hypothesized broader process in the laboratory's MRI experiment. The unique device consists of two concentric cylinders that rotate at different speeds. In this experiment, researchers filled the cylinders with water and attached a water-filled plastic ball tethered by a spring to a post in the center of the device; the stretching and bending spring mimicked the magnetic forces in the plasma in accretion disks. Researchers then rotated the cylinders and videoed the behavior of the ball as seen from the top down.



Simulated accretion disk swirling around a celestial body. Credit: Michael Owen and John Blondin, North Carolina State University.

The findings, reported in *Communications Physics*, compared the motions of the spring-tethered ball when rotating at different speeds. "With no stretching, nothing happens to the angular momentum," said Hantao Ji, a professor of astrophysical sciences at Princeton University and principal researcher on the MRI and a coauthor of the paper. "Nothing also happens if the spring is too strong."

However, direct measurement of the results found that when the spring-

tethering was weak—analogous to the condition of the magnetic fields in accretion disks —behavior of the angular momentum of the ball was consistent with MRI predictions of developments in a real accretion disk. The findings showed that the weakly tethered rotating ball gained angular momentum and shifted outward during the experiment. Since the [angular momentum](#) of a rotating body must be conserved, any gains in momentum must be matched by a loss of momentum in the inner section, allowing gravity to draw the disk into the object it has been orbiting.

**More information:** Derek M. H. Hung et al, Experimental confirmation of the standard magnetorotational instability mechanism with a spring-mass analogue, *Communications Physics* (2019). [DOI: 10.1038/s42005-018-0103-7](#)

Provided by Princeton Plasma Physics Laboratory

Citation: Novel experiment validates widely speculated mechanism behind the formation of stars (2019, February 5) retrieved 9 April 2024 from <https://phys.org/news/2019-02-validates-widely-speculated-mechanism-formation.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.