

New research from Arctic: Thawing permafrost peatlands may add to atmospheric CO₂ burden

March 1 2019




Sampling site near Kilpisjärvi, Finnish Lapland. Credit: Carolina Voigt

A new study led by researchers from the University of Eastern Finland and the University of Montreal, in cooperation with researchers from various Nordic research institutions, finds that peatlands may strengthen the permafrost-carbon feedback by adding to the atmospheric CO₂ burden post-thaw.

Temperatures in the Arctic are rising twice as fast as in the rest of the world, causing [permafrost soils](#) to thaw. Permafrost peatlands are biogeochemical hot spots in the Arctic as they store vast amounts of carbon. Permafrost thaw could release part of these long-term immobile carbon stocks as the greenhouse gases carbon dioxide (CO₂) and methane (CH₄) to the atmosphere, but how much, at which time-span and as which gaseous carbon species is still highly uncertain.

A new study led by researchers from the University of Eastern Finland and the University of Montreal, in cooperation with researchers from various Nordic research institutions, finds that peatlands may strengthen the [permafrost](#)-carbon feedback by adding to the atmospheric CO₂ burden post-thaw. The study was recently published in *Global Change Biology* - a leading journal in [environmental science](#).

By applying a novel experimental approach using intact plant-soil systems (mesocosms), the authors were able to simulate permafrost thaw (thawing of the upper 10-15cm of permafrost) under near-[natural conditions](#). Greenhouse gas flux dynamics were monitored via high-resolution flowthrough gas measurements, combined with detailed monitoring of soil greenhouse gas concentration dynamics, yielding insights into gas production and consumption potential of individual soil layers. The study finds that under dry conditions, peatlands may strengthen the permafrost-carbon feedback by adding to the atmospheric CO₂ burden post-thaw. However, as long as the [water table](#) remains low, the results reveal a strong CH₄ sink capacity in these types of Arctic ecosystems pre- and post-thaw, with the potential to compensate for part of the permafrost CO₂ losses over longer timescales.



Intact plant-soil systems (peat mesocosms) set up in a climate-controlled chamber. Credit: Carolina Voigt

More information: Carolina Voigt et al, Ecosystem carbon response

of an Arctic peatland to simulated permafrost thaw, *Global Change Biology* (2019). DOI: [10.1111/gcb.14574](https://doi.org/10.1111/gcb.14574)

Provided by University of Eastern Finland

Citation: New research from Arctic: Thawing permafrost peatlands may add to atmospheric CO2 burden (2019, March 1) retrieved 22 March 2023 from <https://phys.org/news/2019-03-arctic-permafrost-peatlands-atmospheric-co2.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.