


Race at the edge of the sun: Ions are faster than atoms

March 26 2019



Solar prominence taken from SOT (Solar Optical Telescope) onboard the HINODE satellite showing approx 150,000 km of the curved edge of the sun
Credit: Hinode JAXA/NASA


Scientists at the University of Göttingen, the Institut d'Astrophysique in

Paris and the Istituto Ricerche Solari Locarno have observed that ions move faster than atoms in the gas streams of a solar prominence. The results of their study were published in *The Astrophysical Journal*.

In astrophysics, the "fourth state" of matter plays a crucial role. Apart from solid, liquid and gaseous states, there is also plasma, an accumulation of atoms that have lost shell electrons through collisions or high-energy radiation, and thus become ions. These ions are subject to magnetic forces that do not affect electrically [neutral atoms](#). If there are not too many collisions in the plasma, both particle types can flow independently of each other.

The researchers have now succeeded in observing the physics phenomena in just such a partially ionised plasma without impact equilibrium in gas streams of the sun. The result: in clouds above the edge of the sun, also known as prominences, ions of the element strontium move 22 percent faster than sodium atoms.

Sixteen hours later, the ions were only 11 percent faster. "Evidently, the neutral sodium atoms were more strongly carried along by the strontium ions," says Dr. Eberhard Wiehr of the University of Göttingen, first author of the study. This could be caused by an increased particle density, which increases the probability of impact. "In addition, the flow behaviour of the prominence could have changed in the 16 hours," says Wiehr.



Solar prominence measuring approx 50,000 km above the solar edge taken at the Tenerife Observatory Credit: Dr Eberhard Wiehr

The faster ions move in sync with the oscillation of the magnetic fields. This keeps the prominence in suspension despite the attraction of the sun. Movements in deeper layers of the sun cause the magnetic lines of force to fluctuate. The ions immediately follow the reversal of the direction of oscillation, while the neutral [atoms](#) have to repeatedly

reorient themselves with the ions.

The researchers are now planning a systematic search for prominences with suitable oscillations that can be measured over a longer period of time.

More information: E. Wiehr et al. Evidence for the Two-fluid Scenario in Solar Prominences, *The Astrophysical Journal* (2019). [DOI: 10.3847/1538-4357/ab04a4](https://doi.org/10.3847/1538-4357/ab04a4)

Provided by University of Göttingen

Citation: Race at the edge of the sun: Ions are faster than atoms (2019, March 26) retrieved 12 April 2024 from <https://phys.org/news/2019-03-edge-sun-ions-faster-atoms.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.