

Geologic evidence supports theory that major cosmic impact event occurred approximately 12,800 years ago

March 13 2019, by Sonia Fernandez



The researchers found evidence of cosmic impact at the Pilauco dig site in a suburb of the Osorno province in Chile Credit: Courtesy image

When UC Santa Barbara geology professor emeritus James Kennett and colleagues set out years ago to examine signs of a major cosmic impact

that occurred toward the end of the Pleistocene epoch, little did they know just how far-reaching the projected climatic effect would be.

"It's much more extreme than I ever thought when I started this work," Kennett noted. "The more work that has been done, the more extreme it seems."

He's talking about the Younger Dryas Impact Hypothesis, which postulates that a fragmented comet slammed into the Earth close to 12,800 years ago, causing rapid climatic changes, megafaunal extinctions, sudden human population decrease and cultural shifts and widespread wildfires (biomass burning). The hypothesis suggests a possible triggering mechanism for the abrupt changes in climate at that time, in particular a rapid cooling in the Northern Hemisphere, called the Younger Dryas, amid a general global trend of natural warming and ice sheet melting evidenced by changes in the fossil and sediment record.

Controversial from the time it was proposed, the hypothesis even now continues to be contested by those who prefer to attribute the end-Pleistocene reversal in warming entirely to terrestrial causes. But Kennett and fellow stalwarts of the Younger Dryas Boundary (YDB) Impact Hypothesis, as it is also known, have recently received a major boost: the discovery of a very young, 31-kilometer-wide impact crater beneath the Greenland ice sheet, which they believe may have been one of the many comet fragments that impacted Earth at the onset of the Younger Dryas.

Now, in a paper published in the journal *Nature Scientific Reports*, Kennett and colleagues, led by Chilean paleontologist Mario Pino, present further evidence of a cosmic impact, this time far south of the equator, that likely lead to biomass burning, climate change and megafaunal extinctions nearly 13,000 years ago.

"We have identified the YDB layer at high latitudes in the Southern Hemisphere at near 41 degrees south, close to the tip of South America," Kennett said. This is a major expansion of the extent of the YDB event." The vast majority of evidence to date, he added, has been found in the Northern Hemisphere.

This discovery began several years ago, according to Kennett, when a group of Chilean scientists studying sediment layers at a well-known Quaternary paleontological and [archaeological site](#), Pilauco Bajo, recognized changes known to be associated with YDB impact event. They included a "black mat" layer, 12,800 years in age, that coincided with the disappearance of South American Pleistocene megafauna fossils, an abrupt shift in regional vegetation and a disappearance of human artifacts.

"Because the sequencing of these events looked like what had already been described in the YDB papers for North America and Western Europe, the group decided to run analyses of impact-related proxies in search of the YDB layer," Kennett said. This yielded the presence of microscopic spherules interpreted to have been formed by melting due to the extremely high temperatures associated with impact. The layer containing these spherules also show peak concentrations of platinum and gold, and native iron particles rarely found in nature.

"Among the most important spherules are those that are chromium-rich," Kennett explained. The Pilauco site spherules contain an unusual level of chromium, an element not found in Northern Hemisphere YDB impact spherules, but in South America. "It turns out that volcanic rocks in the southern Andes can be rich in chromium, and these rocks provided a local source for this chromium," he added. "Thus, the cometary objects must have hit South America as well."

Other evidence, which, Kennett noted, is consistent with previous and

ongoing documentation of the region by Chilean scientists, pointed to a "very large environmental disruption at about 40 degrees south." These included a large biomass burning event evidenced by, among other things, micro-charcoal and signs of burning in pollen samples collected at the impact layer. "It's by far the biggest burn event in this region we see in the record that spans thousands of years," Kennett said. Furthermore, he went on, the burning coincides with the timing of major YDB-related burning events in North America and western Europe.

The [sedimentary layers](#) at Pilauco contain a valuable record of pollen and seeds that show change in character of regional vegetation—evidence of a shifting climate. However, in contrast to the Northern Hemisphere, where conditions became colder and wetter at the onset of the Younger Dryas, the opposite occurred in the Southern Hemisphere.

"The plant assemblages indicate that there was an abrupt and major shift in the vegetation from wet, cold conditions at Pilauco to warm, dry conditions," Kennett said. According to him, the atmospheric zonal climatic belts shifted "like a seesaw," with a synergistic mechanism, bringing warming to the Southern Hemisphere even as the Northern Hemisphere experienced cooling and expanding sea ice. The rapidity—within a few years—with which the climate shifted is best attributed to impact-related shifts in atmospheric systems, rather than to the slower oceanic processes, Kennett said.

Meanwhile, the impact with its associated major environmental effects, including burning, is thought to have contributed to the extinction of local South American Pleistocene megafauna—including giant ground sloths, sabretooth cats, mammoths and elephant-like gomphotheres—as well as the termination of the culture similar to the Clovis culture in the north, he added. The amount of bones, artifacts and megafauna-associated fungi that were relatively abundant in the soil at the Pilauco

site declined precipitously at the impact layer, indicating a major local disruption.

The distance of this recently identified YDB site—about 6,000 kilometers from the closest well-studied site in South America—and its correlation with the many Northern Hemispheric sites "greatly expands the extent of the YDB impact event," Kennett said. The sedimentary and paleo-vegetative evidence gathered at the Pilauco site is in line with previous, separate studies conducted by Chilean scientists that indicate a widespread burn and sudden major climate shifts in the region at about YDB onset. This new study further bolsters the hypothesis that a cosmic impact triggered the atmospheric and oceanic conditions of the Younger Dryas, he said.

"This is further evidence that the Younger Dryas climatic onset is an extreme global event, with major consequences on the animal life and the human life at the time," Kennett said. "And this Pilauco section is consistent with that."

More information: Mario Pino et al, Sedimentary record from Patagonia, southern Chile supports cosmic-impact triggering of biomass burning, climate change, and megafaunal extinctions at 12.8 ka, *Scientific Reports* (2019). [DOI: 10.1038/s41598-018-38089-y](https://doi.org/10.1038/s41598-018-38089-y)

Provided by University of California - Santa Barbara

Citation: Geologic evidence supports theory that major cosmic impact event occurred approximately 12,800 years ago (2019, March 13) retrieved 19 April 2024 from <https://phys.org/news/2019-03-geologic-evidence-theory-major-cosmic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.