

Study identifies a key to soybean cyst nematode growth

March 28 2019



How do female soybean cyst nematodes change shape to accommodate their huge reproductive output? New research from the University of Illinois highlights the role of stem-cell-like seam cells. Credit: Nathan Schroeder, University of Illinois

The soybean cyst nematode, one of the crop's most destructive pests, isn't like most of its wormy relatives. Whereas the vast majority of nematodes look like the microscopic worms they are, the female soybean cyst nematode shape-shifts into a tiny lemon after feeding on soybean roots. In a new *EvoDevo* article, a University of Illinois research



team explains how it happens and why.

"We think the soybean cyst <u>nematode</u> has evolved this body shape so that they can produce a lot more offspring," says Nathan Schroeder, assistant professor in the Department of Crop Sciences at the University of Illinois and corresponding author on the new study. "If you compare the most closely <u>related species</u> that stay long and skinny, they have a lot fewer babies than this lady does."

The round shape allows female soybean cyst nematodes to retain about two-thirds of their fertilized eggs inside their bodies. As embryos develop, the mother's body hardens to become a protective cyst. Schroeder says these adaptations have allowed the soybean cyst nematode to become as successful as it is.

From previous research with a different species, the scientists suspected seam cells were responsible for the shift from long and skinny to fat and round. Seam cells, which have stem-cell-like properties, run the length of these worms and divide to enlarge the epidermis every time the worms molt.

"Normally, there's one division for each molt, creating one new set of nuclei in the multinucleated epidermis. We found that with <u>soybean cyst</u> <u>nematode</u>, they divide multiple times after infection, molt, divide even more times, molt, divide even more times. You have <u>exponential growth</u> which leads to this fat, round beast that has lots and lots of these epidermal nuclei," Schroeder explains. "Between the last juvenile phase, which is long and skinny, and the reproductive adult phase, the number of nuclei increase forty-fold."

The research team also investigated the division pattern in other plantparasitic nematodes and found similar seam cell proliferation in several others, despite not being closely related to <u>soybean</u> cyst nematodes.



Essentially, they found evidence of convergent evolution, or the appearance of similar traits to meet the same needs in distantly related species. The common example is wings in butterflies and bats, but now nematode <u>body shape</u> can be added to the list.

"The larger question is how do you get new body shapes in biology? How do new shapes arise? In this case, we're suggesting that these organisms are using mechanisms of stem cell proliferation to generate new <u>body</u> shapes, which is pretty interesting," Schroeder says.

The discovery could also have potential management implications down the road. If researchers could disrupt the proliferation of these seam cells at a certain stage, it might be possible to keep female nematodes from becoming quite so round, with room for so many babies.

"From my perspective, it's a combination of interesting biology and practical implications with an economically important pest causing yield loss throughout the Midwest," Schroeder says. "If we can find some new target strategies that will affect it but not the other thousands of beneficial nematode species in the soil, that would be a game-changer."

More information: Sita Thapa et al, Convergent evolution of saccate body shapes in nematodes through distinct developmental mechanisms, *EvoDevo* (2019). <u>DOI: 10.1186/s13227-019-0118-5</u>

Provided by University of Illinois at Urbana-Champaign

Citation: Study identifies a key to soybean cyst nematode growth (2019, March 28) retrieved 23 June 2024 from <u>https://phys.org/news/2019-03-key-soybean-cyst-nematode-growth.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.