


# Living together: How legume roots accommodate two distinct microbial partners

March 1 2019



The effect of the *lan* mutation on root-hair infection threads formation and on early nodule development. Credit: University of Tsukuba

Legumes such as peas and beans form intimate and mutually beneficial partnerships (symbioses) with nitrogen-fixing bacteria, rhizobia. The plant benefits from an enhanced supply of nitrogen, 'fixed' from the air by the rhizobia, while the bacteria benefit from protective accommodation inside special structures, called root nodules, that supply nutrients from the host plant. A different type of symbiosis is formed between the roots of many plant species and soil fungi, called mycorrhizal fungi. Both types of complex plant-microbe interactions are crucial for supplying plants with nutrients, but many details of how these symbioses develop remain unclear.

University of Tsukuba researchers, collaborating with two other Japanese universities, have revealed a key piece in the jigsaw puzzle of mechanisms that control the developmental processes behind the symbioses of roots with microbes. The team identified a gene that is pivotal in controlling how legume roots establish cellular accommodation for rhizobia bacteria, described in a recent publication in *PLOS Genetics*.

When the gene is inactivated in a mutant plant, the roots produce dramatically fewer nitrogen-fixing nodules because the usual (intracellular) route of entry for the bacteria, called an infection thread, does not develop properly in the root cells. However, small numbers of nodules do develop, some weeks later than normal, when the bacteria enter by a different (intercellular) route between the root cells.

The team named the gene lack of symbiont accommodation (*lan*). They showed it is inactivated in the mutant, and found it is closely related to a gene in other plant species. It is thought to encode a protein that acts in a complex of other regulatory proteins (called Mediator) to control the expression of numerous [genes](#) and processes. This is the first report of the gene's involvement in controlling plant-microbe interactions.

"We used the model legume *Lotus japonicus*, which grows and reproduces rapidly and has a smaller, simpler genome than most crop plants," says corresponding author Takuya Suzuki. "Our [research methods](#) included [genetic modification](#), studying the plant's anatomy by microscopy using [fluorescent dyes](#), genome sequencing, and producing mutant plants using the latest gene editing technologies."

The *lan* gene is important not only for symbiosis with rhizobia: the team showed that the gene is also required for establishing symbioses with mycorrhizal fungi.

"This study shows that a single control system operates in establishing

two completely different symbioses that are important for plant nutrition," says Suzaki. "Our results have wider implications for understanding how plant developmental processes are coordinated."

**More information:** Takuya Suzaki et al, Lack of symbiont accommodation controls intracellular symbiont accommodation in root nodule and arbuscular mycorrhizal symbiosis in *Lotus japonicus*, *PLOS Genetics* (2019). [DOI: 10.1371/journal.pgen.1007865](https://doi.org/10.1371/journal.pgen.1007865)

Provided by University of Tsukuba

Citation: Living together: How legume roots accommodate two distinct microbial partners (2019, March 1) retrieved 9 April 2024 from <https://phys.org/news/2019-03-legume-roots-accommodate-distinct-microbial.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--