


A trap for positrons

March 1 2019



Dr. Eve Stenson uses spare parts to demonstrate the structure of the positron trap: in the center is the permanent magnet. The wire on the left represents a probe that can be inserted into the trap. It allows researchers to determine the quantity of injected particles successfully captured inside the magnetic field. Credit: Axel Griesch / IPP

For the first time, scientists from the Technical University of Munich



(TUM) and the Max Planck Institute for Plasma Physics (IPP) have succeeded in losslessly guiding positrons, the antiparticles of electrons, into a magnetic field trap. This is an important step toward creating a matter-antimatter plasma of electrons and positrons like the plasmas believed to occur near neutron stars and black holes. In an interview, Dr. Eve Stenson presents her research work.

Why do you want to lure positrons into a trap?

Being able to capture and confine positrons is fundamental for studying what is known as electron-positron pair <u>plasma</u>. Such plasmas are of great interest both for the investigation of fundamental questions in <u>plasma physics</u> as well as in astrophysics.

What is so difficult about catching positrons?

Positrons are the antiparticles of electrons, they have the same properties except that they are positively instead of negatively charged. When a positron hits an electron, both instantly annihilate in a flash of light. And since there are electrons in abundance everywhere on Earth, it is extremely difficult to store positrons in such a way that they survive for at least a while.

Fortunately, we have the most powerful positron source in the world, NEPOMUC (neutron induced positron source Munich), here in Garching, north of Munich, at the Research-Neutronsource Heinz Maier-Leibnitz (FRM II). It can produce 900 million positrons per second.

Plasma physicists have been simulating this electron-positron plasma for 40 years. You have now come a decisive step closer to achieving it in practice. How did you do that?



It is actually very difficult to guide charged particles such as the positive positrons into a magnetic trap. The same rules of <u>physics</u> that confine the particles inside this trap unfortunately also keep out the particles that are supposed to enter.

Our trap has a magnetic field very similar to that of the Earth or other celestial bodies. We came up with the idea of briefly applying an electrical voltage to the edge of the trap to guide the positrons through the magnetic "bars". When we then switch the voltage off again, the positrons remain trapped in the cage. It worked so well, even we were surprised.

How long have you been able to confine the positrons?

... for a little more than a second. No group in the world has yet succeeded in doing this with antimatter in this type of trap.

What are the benefits of the results for plasma physics or other areas?

The aim of the APEX (A Positron-Electron Experiment) Group at the Max Planck Institute for Plasma Physics is to produce a matterantimatter plasma of <u>electrons</u> and positrons and to confine that plasma in a magnetic cage. The first step, however, is to be able to produce and store enough positrons. The next step is to actually create and examine such plasmas.

Astrophysics assume that such exotic plasmas occur in the vicinity of <u>neutron stars</u> and black holes. In terrestrial plasma physics, the symmetry of <u>positron</u> and electron masses is expected to lead to new findings on waves and turbulence in plasmas—findings that could help us to use nuclear fusion for power generation in the future.



More information: J. Horn-Stanja et al, Confinement of Positrons Exceeding 1 s in a Supported Magnetic Dipole Trap, *Physical Review Letters* (2018). DOI: 10.1103/PhysRevLett.121.235003

Provided by Technical University Munich

Citation: A trap for positrons (2019, March 1) retrieved 31 August 2024 from <u>https://phys.org/news/2019-03-positrons.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.